

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
6 September 2002 (06.09.2002)

PCT

(10) International Publication Number  
**WO 02/068380 A1**

(51) International Patent Classification<sup>7</sup>: **C07C 275/18**,  
275/34, 335/16, C07D 333/38, C07C 233/52, 233/81,  
237/38, 235/54, 233/63, 235/62, C07D 209/42, 217/26,  
213/81, 213/82, 241/44, C07C 235/84, C07D 233/90,  
C07C 229/48, 271/38, 237/14, C07D 263/56, 277/64,  
A61K 31/00, A61P 25/18, 25/30, 13/04, 25/00

[US/US]; 812 Eaglewood Drive, Zionsville, IN 46077 (US).

(21) International Application Number: PCT/US02/01247

(22) International Filing Date: 12 February 2002 (12.02.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/270,512 22 February 2001 (22.02.2001) US  
60/340,607 30 October 2001 (30.10.2001) US

(74) Agents: **ANDERSON, Arvie, J.** et al.; Eli Lilly and Company, Lilly Corporate Center, Indianapolis, IN 46285 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT (utility model), AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ (utility model), DE (utility model), DK (utility model), DM, DZ, EC, EE (utility model), ES, FI (utility model), GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK (utility model), SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.

(71) Applicant (*for all designated States except US*): **ELI LILLY AND COMPANY** [US/US]; Lilly Corporate Center, Indianapolis, IN 46285 (US).

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **MASSEY, Steven, Marc** [US/US]; 3525 Waterstone Court, Indianapolis, IN 46268 (US). **MONN, James, Allen** [US/US]; 7518 Allenwood Court, Indianapolis, IN 46268 (US). **PRIETO, Lourdes** [ES/ES]; Lilly, S. A., Avda. Industria, 30, 28100-Alcobenda Madrid (ES). **VALLI, Matthew, John**

Published:

— with international search report

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: SYNTHETIC EXCITATORY AMINO ACIDS

(57) Abstract: The present invention provides novel compounds that affect certain excitatory amino acid receptors, and are useful in the treatment of neurological disorders and psychiatric disorders.



**WO 02/068380 A1**

## SYNTHETIC EXCITATORY AMINO ACIDS

## Background of the Invention

In the mammalian central nervous system (CNS), the transmission of nerve impulses is controlled by the interaction between a neurotransmitter, that is released by a sending neuron, and a surface receptor on a receiving neuron, causing excitation of this receiving neuron. L-Glutamate, which is the most abundant neurotransmitter in the CNS, mediates the major excitatory pathway in mammals, and is referred to as an excitatory amino acid (EAA). The receptors that respond to glutamate are called excitatory amino acid receptors (EAA receptors). See Watkins & Evans, *Ann. Rev. Pharmacol. Toxicol.*, 21, 165 (1981); Monaghan, Bridges, and Cotman, *Ann. Rev. Pharmacol. Toxicol.*, 29, 365 (1989); Watkins, Krogsgaard-Larsen, and Honore, *Trans. Pharm. Sci.*, 11, 25 (1990). The excitatory amino acids are of great physiological importance, playing a role in a variety of physiological processes, such as long-term potentiation (learning and memory), the development of synaptic plasticity, motor control, respiration, cardiovascular regulation, emotional states and sensory perception.

The excessive or inappropriate stimulation of excitatory amino acid receptors leads to neuronal cell damage or loss by way of a mechanism known as excitotoxicity. This process has been suggested to mediate neuronal degeneration in a variety of conditions. The medical consequences of such neuronal degeneration make the abatement of these degenerative neurological processes an important therapeutic goal.

Excitatory amino acid receptors are classified into two general types. Receptors that are directly coupled to the opening of cation channels in the cell membrane of the neurons are termed "ionotropic." This type of receptor has

been subdivided into at least three subtypes, which are defined by the depolarizing actions of the selective agonists *N*-methyl-D-aspartate (NMDA),  $\alpha$ -amino-3-hydroxy-5-methylisoxazole-4-propionic acid (AMPA), and kainic acid (KA). The second general type of receptor is the G-protein or second messenger-linked "metabotropic" excitatory amino acid receptor. This second type is coupled to multiple second messenger systems that lead to enhanced phosphoinositide hydrolysis, activation of phospholipase D, increases or decreases in cAMP formation, and changes in ion channel function. Schoepp and Conn, *Trends in Pharmacol. Sci.*, 14, 13 (1993). Both types of receptors appear not only to mediate normal synaptic transmission along excitatory pathways, but also participate in the modification of synaptic connections during development and throughout life. Schoepp, Bockaert, and Sladeczek, *Trends in Pharmacol. Sci.*, 11, 508 (1990); McDonald and Johnson, *Brain Research Reviews*, 15, 41 (1990).

The metabotropic glutamate receptors are a highly heterogeneous family of glutamate receptors that are linked to multiple second-messenger pathways. Generally, these receptors function to modulate the presynaptic release of glutamate, and the postsynaptic sensitivity of the neuronal cell to glutamate excitation. The metabotropic glutamate receptors (mGluR) have been pharmacologically divided into two subtypes. One group of receptors is positively coupled to phospholipase C, which causes hydrolysis of cellular phosphoinositides (PI). This first group are termed PI-linked metabotropic glutamate receptors. The second group of receptors is negatively coupled to adenylyl cyclase, which prevents the forskolin-stimulated accumulation of cyclic adenosine monophosphate (cAMP). Schoepp and Conn, *Trends Pharmacol. Sci.*, 14, 13 (1993). Receptors within this second group are termed cAMP-linked metabotropic glutamate receptors. Agonists of the cAMP-linked metabotropic glutamate receptors should be useful for the treatment of

acute and chronic neurological conditions and psychiatric conditions.


Compounds have recently been discovered that affect metabotropic glutamate receptors, but have no effect on ionotropic glutamate receptors. (1*S*,3*R*)-1-Aminocyclopentane-1,3-dicarboxylic acid (1*S*,3*R*-ACPD) is an agonist of PI-linked and cAMP-linked metabotropic glutamate receptors. Schoepp, Johnson, True, and Monn, *Eur. J. Pharmacol.*, 207, 351 (1991); Schoepp, Johnson, and Monn, *J. Neurochem.*, 58, 1184 (1992). (2*S*,3*S*,4*S*)-2-(carboxycyclopropyl)glycine (L-CCG-I) was recently described as a selective cAMP-linked metabotropic glutamate receptor agonist; however, at higher concentrations, this compound has activity at PI-linked metabotropic receptors. Nakagawa, et al., *Eur. J. Pharmacol.*, 184, 205 (1990); Hayashi, et al., *Br. J. Pharmacol.*, 107, 539 (1992); Schoepp et al., *J. Neurochem.*, 63., page 769-772 (1994).

United States Patent No. 5,958,960 discloses that certain 2-aminobicyclo[3.1.0]hexane-2,6-dicarboxylic acid derivatives are modulators of metabotropic glutamate function, in particular agonists or antagonists of glutamate at metabotropic glutamate receptors and as such are useful in the treatment of a neurological disorder or a psychiatric disorder that has been linked to the excitatory amino acid receptors.

#### Summary of the Invention

The present invention provides compounds that selectively affect the negatively coupled cAMP-linked metabotropic glutamate receptors. More specifically, the present invention relates to compounds of the formula

- 4 -



I

wherein:


X is CH<sub>2</sub>, O, or NH;

Y is O, S, N or H;

A is a bond, O, N, (1-10C) alkyl, (2-10C) alkenyl or (2-10C) alkynyl;

R is hydrogen, (1-10C) alkyl, (2-10C) alkenyl, (3-6C) alkynyl, aryl, heterocyclyl or substituted aryl;

or the group XC(Y)AR is



where Q is O, S or NH;

or a pharmaceutically acceptable metabolically labile ester or amide thereof;

or a pharmaceutically acceptable salt thereof.

A particular compound of formula I is one wherein (1-10C) alkyl is methyl, ethyl, n-propyl, isopropyl, n-butyl, t-butyl, n-pentyl, isopentyl, n-hexyl, heptyl, n-octyl, nonyl or decyl; (2-10C) alkenyl is allyl, allenyl, 1-butenyl, 1-pentenyl, 3-nonenyl or 5-decenyl; (2-6C) alkynyl is ethynyl, propynyl, butynyl or pentynyl; aryl is phenyl, substituted phenyl or naphthyl; and arylalkyl is benzyl, 2-nitro benzyl, or 1-phenylethyl.

Another particular compound of formula I is one wherein R is 2-methoxyphenyl, 3-methoxyphenyl, 4-methoxyphenyl, 3-methylphenyl, 4-methylphenyl, 2-

chlorophenyl, 3-chlorophenyl, 4-chlorophenyl, 3-fluorophenyl, 2-fluorophenyl, 4-fluorophenyl, 2,6-dichlorophenyl, 2,4-dichlorophenyl, 3,5-dichlorophenyl, 3,4-dichlorophenyl, methyl, 2-naphthyl, 1-naphthyl, 3-methylphenyl, phenyl, thiophenyl, 3-trifluoromethylphenyl, 2,3-dichlorophenyl, 1H-indole-3-yl cyclopropanyl, 1H-indol-2-yl, 4-hydroxyphenyl, 3-hydroxyphenyl, 2-hydroxyphenyl, 4-biphenyl, 1-isoquinolyl, 3-pyridinyl, 2-pyridinyl, 3,5-difluorophenyl, 4-pyridinyl, 2-methylphenyl, 2-quinoxaliny, hydrogen, 3-carboxyphenyl, 2-trifluoromethylphenyl, benzyl, 4-trifluoromethylphenyl;

A is a bond, methyl, O, NH, ethenyl or ethynyl;

Y is O, S or H,H; and

X is CH<sub>2</sub>, O or NH.

Another particular compound of formula I is one wherein R is heterocyclyl or substituted aryl.

Another particular compound of formula I is one wherein R is 1H-indolyl, 2-naphthyl, 3-chlorophenyl or 2-methoxyphenyl.

Another particular compound of formula I is one wherein Y is O or S.

Another particular compound of formula I is one wherein A is CH<sub>2</sub> or a bond.

A preferred compound of formula I is one wherein (1S\*, 2S\*, 4S\*, 5R\*, 6S\*) 2-amino-4-[3-3-chlorophenyl)ureido]bicyclo[3.1.0]hexane-2,6-dicarboxylic acid.

The compounds of formula I are modulators of metabotropic glutamate receptor function with improved activity, in particular potent agonists of mGluR2 and mGluR3 receptors.


According to another aspect, therefore, the present invention provides a method of modulating metabotropic glutamate receptor function in a mammal including a human, which comprises administering an effective amount of a

compound of formula I, or a non-toxic metabolically labile ester or amide thereof, or a pharmaceutically acceptable salt thereof.

According to yet another aspect, the present invention provides the use of a compound of formula I as defined hereinabove for the manufacture of a medicament for use in modulating metabotropic glutamate receptor function.

The present invention also includes all physical forms of the compounds of formula I, including crystalline solvates.

It will be appreciated that the compounds of formula I contain at least five asymmetric carbon atoms. As used herein, a wedge line refers to a bond projecting from the plain of the paper to the reader's eyes and is referred to as "beta" stereochemistry. A dashed line refers to a bond projecting into the plain of the paper away from the reader's eyes and is referred to as "alpha" stereochemistry. A plain line refers to a bond lying within the plain of the paper. Preferably the stereoisomer of compounds of formula I have the configuration I with functional groups attached at C-4 possessing beta stereochemistry, as shown below.




The present invention also provides pharmaceutical formulations comprising a compound of formula I in combination with one or more pharmaceutically acceptable carriers, diluents, or excipients.

Further aspects of the present invention include a method for affecting the cAMP-linked metabotropic glutamate

receptors, as well as methods for treating a neurological disorder or a psychiatric disorder that has been linked to the excitatory amino acid receptors, which comprises administering a compound of formula I. Examples of neurological disorders that are treated with a formula I compound include cerebral deficits subsequent to cardiac bypass surgery and grafting, cerebral ischemia (e.g. stroke and cardiac arrest); spinal cord trauma; head trauma; Alzheimer's Disease; Huntington's Chorea; amyotrophic lateral sclerosis; AIDS-induced dementia; muscular spasms; migraine headaches; urinary incontinence; convulsions; perinatal hypoxia; hypoglycemic neuronal damage; drug tolerance, withdrawal, and cessation (i.e. opiates, benzodiazepines, nicotine, cocaine, or ethanol); smoking cessation; ocular damage and retinopathy; cognitive disorders; idiopathic and drug-induced Parkinson's Disease; emesis; brain edema; chronic pain; sleep disorders; Tourette's syndrome; attention deficit disorder; and tardive dyskinesia. Examples of psychiatric disorders that are treated with a formula I compound include schizophrenia, anxiety and related disorders (e.g. panic attack and stress-related disorders), depression, bipolar disorders, psychosis, and obsessive compulsive disorders.

The present invention also provides a process for producing a compound of formula I, or a pharmaceutically salt thereof, which comprises:

(1) deprotecting the amine group of a compound of formula



II



wherein:

$R^1$  is a nitrogen protecting group such t-butoxycarbonyl; and

$R^2$  and  $R^3$  are both carboxy protecting groups such as ethyl;

(2) for a compound of formula I deprotecting the carboxy groups of a compound of formula II where  $R^2$  and  $R^3$  are both carboxy protecting groups such as ethyl;

(3) optionally preparing a pharmaceutically-acceptable salt of the formula I compound.

#### Detailed Description of the Invention

The term "(1-10C) alkyl" represents a straight, branched, or cyclic alkyl chain having from one to ten carbon atoms. Typical straight or branched  $C_1$ - $C_{10}$  alkyl groups include methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, t-butyl, n-pentyl, isopentyl, neopentyl, n-hexyl, 2-methylpentyl, 3-methylpentyl, 4-methylpentyl, 2,2-dimethylbutyl, 2,3-dimethylbutyl, 3,3-dimethylbutyl, heptyl, n-octyl, 2,2-dimethylhexyl, 2,5-dimethylhexyl, 2-methylheptyl, 4-methylheptyl, 2,2,4-trimethylpentyl, 2,3,4-trimethylpentyl, nonyl, 3,5,5-trimethylhexyl, decyl, 3,7-dimethyloctyl, and the like. The term "(1-10C) alkyl" includes within it the terms " $C_1$ - $C_6$  alkyl" and " $C_1$ - $C_4$  alkyl". Typical cyclic alkyl groups include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, and the like. Typical  $C_1$ - $C_6$  alkyl groups include methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, t-butyl, n-pentyl, isopentyl, neopentyl, and n-hexyl.

The term "(2-10C) alkenyl" represents straight or branched unsaturated alkyl chains having from two to ten carbon atoms, and having one or more carbon-carbon double bond, such as, dienes and trienes. This group also includes

both E and Z isomers. Representative radicals for this group include ethenyl, allyl, allenyl, 1-butenyl, 2-butenyl, 2-methyl-1-propenyl, 3-butenyl, 2-methyl-2-propenyl, butadienyl, 1-pentenyl, 2-pentenyl, 2-methyl-2-butenyl, 4-pentenyl, 3-methyl-2-butenyl, 3-methyl-1,2-butadienyl, 3-hexenyl, 2-hexenyl, 4-methyl-3-pentenyl, 4-hexenyl, 5-hexenyl, 3-methyl-1-penten-3-yl, 4-methyl-3-pentenyl, 6-methyl-5-heptene-2-yl, 7-octenyl, 1-octen-3-yl, 3-nonenyl, 2,4-dimethyl-2,6-heptadienyl, 3,7-dimethyl-6-octenyl, 5-decenyl, 9-decenyl, 2,6-dimethyl-7-octenyl, and the like. The term "(2-10C) alkenyl" includes within it the term "(2-6C) alkenyl".

The term "(2-6C) alkynyl" represents ethynyl, propynyl, butynyl or pentynyl.

The phrase "stereoisomeric compound" represents an optical isomer of a Formula I compound, which includes the 1S, 2S, 4S, 5R, 6R isomer.

The term "carboxy protecting group" as used herein refers to one of the ester derivatives of the carboxylic acid group commonly employed to block or protect the carboxylic acid group while reactions are carried out on other functional groups. The protection of carboxylic acid groups is generally described in McOmie, Protecting Groups in Organic Chemistry, Plenum Press, NY, 1973; and Greene and Wuts, Protecting Groups in Organic Synthesis, 2nd. Ed., John Wiley & Sons, NY, 1991. Examples of such carboxy protecting groups include methyl, ethyl, methoxymethyl, methylthiomethyl, triphenylmethyl, benzyl, 4-nitrobenzyl, 4-methoxybenzyl, 3,4-dimethoxybenzyl, 2,4-dimethoxybenzyl, 2,4,6-trimethoxybenzyl, 2,4,6-trimethylbenzyl, benzhydryl, t-butyl, t-amyl, trityl, trimethylsilyl, t-butyldimethylsilyl, allyl, 1-(trimethylsilylmethyl)-prop-1-en-3-yl, and the like. Particularly preferred carboxy protecting groups are (C<sub>1</sub>-C<sub>6</sub>) alkyl groups such as ethyl. The term "protected carboxy" refers to a carboxylic acid group having a carboxy protecting group.

The term "nitrogen protecting group" as used herein refers to substituents on amino groups that are commonly employed to block or protect the amino functionality while reactions are carried out in other functional groups. The protection of amino groups is generally described in McOmie, *Protecting Groups in Organic Chemistry*; Plenum Press, NY, 1973, and Greene and Wuts, *Protecting Groups in Organic Synthesis*, 2nd. Ed., John Wiley & Sons, NY, 1991. Examples of nitrogen protecting groups include benzyl, t-butyl, allyl, triphenylmethyl, t-butyldimethylsilyl, triphenylsilyl, formyl, trityl, phthalimido, trichloroacetyl, chloroacetyl, phthaloyl, 2-nitrophenoxyacetyl, benzyloxycarbonyl, methoxycarbonyl, 2-methylbenzyloxycarbonyl, t-butoxycarbonyl, allyloxy-carbonyl, 2,2,2-trichloroethoxycarbonyl, and the like. The term "protected amino" refers to a primary or secondary amine having a nitrogen protecting group.

The term "heterocyclyl" includes heteroaromatics an aromatic 5-6 membered ring containing from one to four heteroatoms selected from oxygen, sulfur and nitrogen, and a bicyclic group consisting of a 5-6 membered ring containing from one to four heteroatoms selected from oxygen, sulfur and nitrogen fused with a benzene ring or a 5-6 membered ring containing from one to four heteroatoms selected from oxygen, sulfur and nitrogen. Examples of heteroaromatic groups are furyl, thiophenyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, imidazolyl, pyrimidyl, benzofuryl, benzothiophenyl, benzimidazolyl, benzoxazolyl, benzo-thiazolyl and indolyl. Examples of particular values are 2-thienyl, 3-thienyl, 2-indolyl, 3-indolyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, 1-isoquinolinyl, 3-isoquinolinyl, 2-benzothiazolyl, benzoxazolyl and 4-imidazolyl.

The term aryl includes phenyl and a polycyclic aromatic carbocyclic ring such as 1-naphthyl or 2-naphthyl.

The term "substituted", as used in the term "substituted heterocycles or aromatic group", herein

signifies that one, two or more substituents may be present, said substituents being selected from atoms and groups which, when present in the compound of formula I, do not prevent the compound of formula I from functioning as a modulator of metabotropic glutamate receptor function.

Examples of atoms and groups which may be present in an optionally substituted heteroaromatic or aryl group are amino, hydroxy, nitro, halogeno, (1-6C) alkyl, (1-6C) alkoxy, (1-6C) alkylthio, carboxy, (1-6C) alkoxycarbonyl, carbamoyl, (1-6C) alkanoylamino, (1-6C) alkylsulphonyl, (1-6C) alkylsulphonylamino, (1-6C) alkanoyl, phenyl, phenoxy, phenylthio, phenylsulphonyl, phenylsulphonylamino, toluene-sulphonylamino, and (1-6C) fluoroalkyl. Examples of particular values are hydroxy, fluoro, chloro, bromo, iodo, methyl, methoxy, carboxy, acetyl, phenyl, phenoxy, tetrazoyl and trifluoromethyl.

Examples of values for substituted aryl groups are 1-naphthyl, 2-naphthyl, phenyl, 2-hydroxyphenyl, 3-hydroxyphenyl, 4-hydroxyphenyl, 3,5-dihydroxyphenyl, 2-fluorophenyl, 3-fluorophenyl, 4-fluorophenyl, 3,5-difluorophenyl, 2-chlorophenyl, 3-chlorophenyl, 4-chlorophenyl, 2,4-dichlorophenyl, 3,4-dichlorophenyl, 3,5-dichlorophenyl, 2-methylphenyl, 3-methylphenyl, 4-methylphenyl, 2-methoxyphenyl, 3-methoxyphenyl, 4-methoxyphenyl, 3-trifluoromethylphenyl, 4-trifluoromethylphenyl, 2-methylphenyl, 3-methylphenyl, 4-methylphenyl, 3-carboxyphenyl, 4-carboxyphenyl and 3-(5'-tetrazoyl)phenyl.

The term heterocyclyl also includes non-aromatic heterocyclyl which includes a 4 to 7 membered ring containing one or two heteroatoms selected from oxygen, sulphur and nitrogen, for example azetidin-1-yl or -2-yl, pyrrolidin-1-yl, -2-yl or -3-yl, piperidin-1-yl, -2-yl, -3-yl or -4-yl, hexahydroazepin-1-yl, -2-yl, -3-yl or -4-yl, oxetan-2-yl or -3-yl, tetrahydro-furan-2-yl or -3-yl, tetrahydropyran-2-yl, -3-yl or -4-yl, hexahydrooxepin-2-yl,

-3-yl or -4-yl, thietan-2-yl or -3-yl, tetrahydrothiophen-2-yl or -3-yl, tetrahydrothiopyran-2-yl, -3-yl or -4-yl, hexahydrothiepin-2-yl, -3-yl or -4-yl, piperazin-1-yl or -2-yl, morpholin-1-yl, -2-yl or -3-yl, thiomorpholin-1-yl, -2-yl or -3-yl, tetrahydropyrimidin-1-yl, -2-yl, -4-yl or -5-yl, imidazolin-1-yl, -2-yl or -4-yl, imidazolidin-1-yl, -2-yl or -4-yl, oxazolin-2-yl, -3-yl, -4-yl or -5-yl, oxazolidin-2-yl, -3-yl, -4-yl or -5-yl, thiazolin-2-yl, -3-yl, -4-yl or -5-yl, or thiazolidin-2-yl, -3-yl, -4-yl or -5-yl.

The term "affecting" refers to a formula I compound acting as an agonist at an excitatory amino acid receptor. The term "excitatory amino acid receptor" refers to a metabotropic glutamate receptor, a receptor that is coupled to cellular effectors via GTP-binding proteins. The term "cAMP-linked metabotropic glutamate receptor" refers to a metabotropic receptor that is coupled to inhibition of adenylate cyclase activity.

The term "neurological disorder" refers to both acute and chronic neurodegenerative conditions, including cerebral deficits subsequent to cardiac bypass surgery and grafting, cerebral ischemia (for example stroke resulting from cardiac arrest), spinal cord trauma, head trauma, Alzheimer's Disease, Huntington's Chorea, amyotrophic lateral sclerosis, AIDS-induced dementia, perinatal hypoxia, hypoglycemic neuronal damage, ocular damage and retinopathy, cognitive disorders, idiopathic and drug-induced Parkinson's Disease. This term also includes other neurological conditions that are caused by glutamate dysfunction, including muscular spasms, migraine headaches, urinary incontinence, drug tolerance, withdrawal, and cessation (i.e. opiates, benzodiazepines, nicotine, cocaine, or ethanol), smoking cessation, emesis, brain edema, chronic pain, sleep disorders, convulsions, Tourette's syndrome, attention deficit disorder, and tardive dyskinesia.

The term "psychiatric disorder" refers to both acute and chronic psychiatric conditions, including schizophrenia, anxiety and related disorders (e.g. panic attack and stress-related cardiovascular disorders), depression, bipolar disorders, psychosis, and obsessive compulsive disorders.

As used herein the term "effective amount" refers to the amount or dose of the compound, upon single or multiple dose administration to the patient, which provides the desired effect in the patient under diagnosis or treatment.

An effective amount can be readily determined by the attending diagnostician, as one skilled in the art, by the use of known techniques and by observing results obtained under analogous circumstances. In determining the effective amount or dose of compound administered, a number of factors are considered by the attending diagnostician, including, but not limited to: the species of mammal; its size, age, and general health; the specific disease involved; the degree of or involvement or the severity of the disease; the response of the individual patient; the particular compound administered; the mode of administration; the bioavailability characteristics of the preparation administered; the dose regimen selected; the use of concomitant medication; and other relevant circumstances. For example, a typical daily dose may contain from about 150 micrograms to about 150 mg of the active ingredient. The compounds can be administered by a variety of routes including oral, rectal, transdermal, subcutaneous, intravenous, intramuscular, buccal or intranasal routes. Alternatively, the compound may be administered by continuous infusion.

As used herein the term "patient" refers to a mammal, such as a mouse, guinea pig, rat, dog or human. It is understood that the preferred patient is a human.

The term "treating" (or "treat") as used herein includes its generally accepted meaning which encompasses prohibiting, preventing, restraining, and slowing, stopping,

or reversing progression of a resultant symptom. As such, the methods of this invention encompass both therapeutic and prophylactic administration.

The present invention includes pharmaceutically-acceptable salts of the formula I compounds. These salts can exist in conjunction with the acidic or basic portion of the molecule and can exist as acid addition, primary, secondary, tertiary, or quaternary ammonium, alkali metal, or alkaline earth metal salts. Generally, the acid addition salts are prepared by the reaction of an acid with a compound of formula I.

Acids commonly employed to form such salts include inorganic acids such as hydrochloric, hydrobromic, hydriodic, sulfuric, and phosphoric acid, as well as organic acids such as *para*-toluenesulfonic, methanesulfonic, oxalic, *para*-bromophenylsulfonic, carbonic, succinic, citric, benzoic, and acetic acid, and related inorganic and organic acids. Such pharmaceutically-acceptable salts thus include sulfate, pyrosulfate, bisulfate, sulfite, bisulfite, phosphate, ammonium, monohydrogenphosphate, dihydrogenphosphate, *meta*-phosphate, pyrophosphate, chloride, bromide, iodide, acetate, propionate, decanoate, caprylate, acrylate, formate, isobutyrate, caprate, heptanoate, propiolate, oxalate, malonate, succinate, suberate, sebacate, fumarate, hippurate, maleate, butyne-1,4-dioate, hexyne-1,6-dioate, benzoate, chlorobenzoate, methylbenzoate, dinitrobenzoate, hydroxybenzoate, methoxybenzoate, phthalate, sulfonate, xylenesulfonate, phenylacetate, phenylpropionate, phenylbutyrate, citrate, lactate,  $\alpha$ -hydroxybutyrate, glycolate, maleate, tartrate, methanesulfonate, propanesulfonate, naphthalene-1-sulfonate, naphthalene-2-sulfonate, mandelate, magnesium, tetramethylammonium, potassium, trimethylammonium, sodium, methylammonium, calcium, and the like salts.

The absolute stereochemical configuration of this most preferred enantiomer has been determined to be 1S, 2S, 4S, 5R, 6S.

While all the formula I compounds of the present invention are believed to selectively affect the negatively-coupled cAMP-linked metabotropic glutamate receptors, certain compounds of the invention are preferred for such use. Preferably, X is NH, O or CH<sub>2</sub>, Y is O, Z is NH or a bond, and R is aryl, heteroaryl, fused aryl or substituted aryl. Representative compounds from this preferred group of formula I compounds include (1S\*, 2S\*, 4S\*, 5R\*, 6S\*) 2-amino-4-(3-chloro-benzoylamino)-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid, (1S\*, 2S\*, 4S\*, 5R\*, 6S\*) 2-amino-4-(3-methoxy-benzoylamino)-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid, (1S\*, 2S\*, 4S\*, 5R\*, 6S\*) 2-amino-4-(3-hydroxy-benzoylamino)-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid, (1S\*, 2S\*, 4S\*, 5R\*, 6S\*) 2-amino-4-(3,4-dichloro-benzoylamino)-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid, (1S\*, 2S\*, 4S\*, 5R\*, 6S\*) 2-amino-4-[(1H-indole-2-carbonyl)-amino]-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid, (1S\*, 2S\*, 4S\*, 5R\*, 6S\*) 2-amino-4-[(naphthalene-1-carbonyl)-amino]-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid, (1S\*, 2S\*, 4S\*, 5R\*, 6S\*) 2-amino-4-[(isoquinoline-1-carbonyl)-amino]-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid, (1S\*, 2S\*, 4S\*, 5R\*, 6S\*)-2-amino-4-(3-carboxy-benzoylamino)-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid, (1S\*, 2S\*, 4S\*, 5R\*, 6S\*) 2-amino-4-[3-(4-fluorophenyl)ureido]bicyclo[3.1.0]hexane-2,6-dicarboxylic acid, (1S\*, 2S\*, 4S\*, 5R\*, 6S\*) 2-amino-4-[3-(3,5-dichlorophenyl)ureido]bicyclo[3.1.0]hexane-2,6-dicarboxylic acid, (1S\*, 2S\*, 4S\*, 5R\*, 6R\*)-2-amino-4-(3-fluorophenylcarbamoyloxy)-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid, and (1S\*, 2S\*, 4R\*, 5R\*, 6S\*) 2-amino-4-(3-chloro)phenylcarbamoylmethyl bicyclo[3.1.0]hexane-2,6-dicarboxylic acid. Certain compounds of the present invention are more preferred for use in affecting the cAMP-



linked metabotropic glutamate receptors. More preferably, R is heteroaromatic, aryl or substituted aryl. Representative compounds from this more preferred group of compounds include (1S\*, 2S\*, 4S\*, 5R\*, 6S\*) 2-amino-4-[(1H-indole-3-carbonyl)-amino]-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid, (1S\*, 2S\*, 4S\*, 5R\*, 6S\*) 2-amino-4-[3-(2-naphthyl)ureido]bicyclo[3.1.0]hexane-2,6-dicarboxylic acid, (1S\*, 2S\*, 4S\*, 5R\*, 6R\*)-2-amino-4-(3-chloro-phenylcarbamoxyloxy)-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid, and (1S\*, 2S\*, 4R\*, 5R\*, 6S\*) 2-amino-4-(2-methoxy)-phenylcarbamoxylmethyl bicyclo[3.1.0]hexane-2,6-dicarboxylic acid.


Most preferably, the compound of formula (I) is (1S\*, 2S\*, 4S\*, 5R\*, 6S\*) 2-amino-4-[3-(3-chlorophenyl)ureido]bicyclo[3.1.0]hexane-2,6-dicarboxylic acid, a C<sub>1</sub>-C<sub>4</sub> alkyl, aralkyl or aryl ester thereof or a pharmaceutically acceptable salt thereof.

While all the compounds of formula II of the present invention are believed to be useful for the synthesis of compounds of formula I, certain compounds are preferred. Preferably, R<sup>1</sup> is t-butoxycarbonyl. More preferably, R<sup>2</sup> and R<sup>3</sup> are C<sub>1</sub>-C<sub>10</sub> alkyl groups for example ethyl groups.


The compounds of formula I of the present invention are generally synthesized from compounds of formula III or formula X where R<sup>1</sup>, R<sup>2</sup> and R<sup>3</sup> are as previously described. The compounds of formula III and formula X are prepared as described in U. S. Patent No. 5,958,960 which is incorporated by reference in its entirety.

Generally, compounds of formula II in which X and Y are O, and A is N may be prepared by directly reacting compounds of formula III. Compounds of formula II in which X and A are N, and Y is O may be prepared by reacting compounds of formula IX. Compounds of formula IX may be prepared by reacting compounds of formula III in a series of steps to convert the hydroxyl to an amine while retaining the stereochemistry at point of attachment.

More specifically, compounds of formula III are reacted with sulphonating agents such as p-toluenesulphonyl chloride in a suitable solvent such as pyridine to produce compounds of formula IV. Compounds of formula IV are reacted with potassium nitrite in DMSO at around 50°C to provide compounds of formula V. It will be appreciated that compounds of formula V have the hydroxyl in opposite configuration to that of the hydroxyl in compounds of formula III.




Compounds of formula V are reacted with sulphonating agents such as p-toluenesulphonyl chloride in a suitable solvent such as pyridine to produce compounds of formula VI. Compounds of formula VI are reacted with an azide salt such as sodium azide for example in dimethylsulfoxide as a reaction solvent to produce compounds of formula VII.




Alternatively, compounds of formula VII may be prepared by reacting compounds of formula III with halogenating agents such as bromine and triphenyl phosphine in the presence of a suitable base such as pyridine in a reaction solvent such as methylene chloride to produce compounds of formula VIII. Compounds of formula VIII are reacted with an azide salt such as sodium azide for example in dimethylsulfoxide as a reaction solvent to produce compounds of formula VII.

-19-



The compounds of formula IX may be prepared by reducing a corresponding compound of formula VII. The reduction is conveniently performed using triphenylphosphine in a suitable solvent such as tetrahydrofuran at a temperature in the range of 0 to 100°C.



Compounds of formula II in which X and A are N, and Y is O may be prepared by reacting compounds of formula IX

with an isocyanate of formula  $R-N=C=O$ . Convenient solvents include dichloromethane.

Compounds of formula II in which X and Y are O, and A is N may be prepared by reacting compounds of formula III with an isocyanate of formula  $R-N=C=O$ . Convenient solvents include dichloromethane.

Compounds of formula II in which X is N, Y is O and A is a bond may be prepared by reacting compounds of formula IX, an amine, with an acyl halide of formula  $RCOX^1$  in which  $X^1$  is, for example, chlorine or bromine. The reaction is conveniently performed in the presence of a base, such as diisopropylethyl amine and in a solvent such as dichloromethane. Alternatively, the amine may be reacted with an acyl isourea such as obtained by the reaction of a carboxylic acid of the formula  $RCOOH$  with a carbodiimide such as dicyclohexylcarbodiimide. The reaction is conveniently performed in the presence of an activating agent such as hydroxybenzotriazole and in a solvent such as dichloromethane.

Generally, compounds of formula II in which X is  $CH_2$ , Y is O, and A is N may be prepared by reacting, in a series of steps, compounds of formula X.

More specifically, compounds of formula XI where  $R^4$  is not hydrogen (a carboxy protecting group) may be prepared from compounds of formula X by a Wadsworth-Emmons reaction, for example by reaction with an alkali metal salt of a dialkyl phosphono acetate ester, such as the sodium salt of allyl diethylphosphonoacetate. The reaction is conveniently performed in an anhydrous solvent such as anhydrous THF. It is appreciated that compounds of formula XI may exist in the (E) or (Z) isomeric form or as a mixture of (E) and (Z) isomers and, as such, are included in the present invention.


Compounds of formula XI where  $R^4$  is hydrogen (a acid) may be prepared from compounds of formula XI where  $R^4$  is not hydrogen (a carboxy protecting group) by procedures well known in the art. In the case where  $R^4$  is allyl, compounds

of formula XI are reacted with a metal catalyst such as chlorotristriphenylphosphine rhodium(I) or tetrakistriphenylphosphine palladium(0) to produce compounds of formula XI where  $R^4$  is hydrogen. The reaction is conveniently performed in the presence of a base such as pyrrolidine and in solvents such as ethanol/water or dichloromethane.

Compounds of formula XII may be prepared by reacting compounds of formula XI where  $R^4$  is hydrogen with oxalyl chloride and catalytic DMF in a suitable solvent such as dichloromethane to produce an intermediate acyl chloride. The intermediate acyl chlorides are reacted with excess amine such as aniline in a suitable solvent such as dichloromethane.

Compounds of formula II where X is  $CH_2$  may be prepared by reacting compounds of formula XII with hydrogen in the presence of a suitable metal catalyst such as palladium on carbon. Convenient solvents include ethanol.

-22-



The formula I compounds of the present invention are agonists of certain metabotropic excitatory amino acid receptors. Specifically, the formula I compounds are agonists of the negatively-coupled cAMP-linked metabotropic glutamate receptors. Therefore, another aspect of the present invention is a method of affecting an excitatory amino acid receptor in mammals, which comprises

administering to a mammal requiring modulated excitatory amino acid neurotransmission a pharmaceutically-effective amount of a compound of formula I. The term "pharmaceutically-effective amount" is used to represent an amount of the compound of the invention which is capable of affecting the excitatory amino acid receptors. By affecting, a compound of the invention is acting as an agonist. When a compound of the invention acts as an agonist, the interaction of the compound with the EAA receptor mimics the response of the interaction of this receptor with its natural ligand (i.e. L-glutamate).

The particular dose of compound administered according to this invention will of course be determined by the particular circumstances surrounding the case, including the compound administered, the route of administration, the particular condition being treated, and similar considerations. The compounds can be administered by a variety of routes including oral, rectal, transdermal, subcutaneous, intravenous, intramuscular, or intranasal routes. Alternatively, the compound may be administered by continuous infusion.

A variety of physiological functions have been shown to be subject to influence by excessive or inappropriate stimulation of excitatory amino acid transmission. The formula I compounds of the present invention are believed to have the ability to treat a variety of neurological disorders in mammals associated with this condition, including acute neurological disorders such as cerebral deficits subsequent to cardiac bypass surgery and grafting, cerebral ischemia (e.g. stroke and cardiac arrest), spinal cord trauma, head trauma, perinatal hypoxia, and hypoglycemic neuronal damage. The formula I compounds are believed to have the ability to treat a variety of chronic neurological disorders, such as Alzheimer's disease, Huntington's Chorea, amyotrophic lateral sclerosis, AIDS-induced dementia, ocular damage and retinopathy, cognitive



disorders, and idiopathic and drug-induced Parkinson's. The present invention also provides methods for treating these disorders which comprises administering to a patient in need thereof an effective amount of a compound of formula I.

The formula I compounds of the present invention are also believed to have the ability to treat a variety of other neurological disorders in mammals that are associated with glutamate dysfunction, including muscular spasms; convulsions; migraine headaches; urinary incontinence; psychosis; drug tolerance, withdrawal, and cessation (i.e. opiates, benzodiazepines, nicotine, cocaine, or ethanol); smoking cessation; anxiety and related disorders (e.g. panic attack and stress-related disorders); emesis; brain edema; chronic pain; sleep disorders; Tourette's syndrome; attention deficit disorder; and tardive dyskinesia. Therefore, the present invention also provides methods for treating these disorders which comprise administering to a patient in need thereof an effective amount of the compound of formula I.

The compounds of the present invention are agonists of cAMP-linked metabotropic glutamate receptors. These compounds are negatively coupled through the receptor to adenylyl cyclase, inhibiting the formation of cyclic adenosine monophosphate. The formula I compounds of the present invention are, therefore, believed to have the ability to treat a variety of psychiatric disorders, such as schizophrenia, anxiety and related disorders (e.g. panic attack and stress-related disorders), depression, bipolar disorders, psychosis, and obsessive compulsive disorders. The present invention also provides methods for treating these disorders which comprises administering to a patient in need thereof an effective amount of a compound of formula I.

To study the ability to affect receptor binding of compounds of the present invention in comparison to compounds of United States Patent No. 5,958,960,

displacement of a high affinity mGluR2 antagonist radioligand [<sup>3</sup>H]LY341495 from recombinant human mGluR2 and human mGluR3 receptors expressed in RGT cells was determined. (See, Ornstein P. L., Arnold M. B., Bleisch T. J., Wright R. A., Wheeler W. J., and Schoepp D. D., [<sup>3</sup>H]LY341495, a highly potent, selective and novel radioligand for labeling group II metabotropic receptors. *Bioorg. Med. Chem. Lett.* 8: 1919-1922 (1998); and Johnson B. G., Wright R. A., Arnold M. B., Wheeler W. J., Ornstein P. L., and Schoepp D. D., [<sup>3</sup>H]LY341495 as a novel rapid filtration antagonist radioligand for group II metabotropic receptors: Characterization of binding to membranes of mGlu receptor subtype expressing cells. *Neuropharmacology* 38: 1519-1529 (1999)). As shown in table 1, compounds of the current invention have improved affinity for the mGlu3 receptors when compared to compounds of United States Patent No. 5,958,960.

Table 1. Effects on [<sup>3</sup>H]LY341495 binding to mGlu2 and mGlu3 receptors.

Compounds of U.S. 5,958,960	K <sub>i</sub> (nM) $\pm$ SEM	
	mGlu2	mGlu3
(1S*, 2S*, 4R*, 5R*, 6S*) 2-Amino-4-(3-methylureido)bicyclo[3.1.0]hexane-2,6-dicarboxylic acid	468.3 $\pm$ 29.0	408.0 $\pm$ 25.4
(1S*, 2S*, 4R*, 5R*, 6S*) 4-Acetylamino-2-aminobicyclo[3.1.0]hexane-2,6-dicarboxylic acid	638.1 $\pm$ 59.6	399.3 $\pm$ 67.8
1S*, 2S*, 4R*, 5R*, 6S*) 2-Amino-4-benzoylamino-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid	607.3 $\pm$ 10.2	393.4 $\pm$ 17.8
Compounds of present invention		
(1S*, 2S*, 4S*, 5R*, 6S*) 2-Amino-4-(3-methylureido)bicyclo[3.1.0]hexane-2,6-dicarboxylic acid	254.2 $\pm$ 10.7	23.0 $\pm$ 2.6
(1S*, 2S*, 4S*, 5R*, 6S*) 4-Acetylamino-2-aminobicyclo[3.1.0]hexane-2,6-dicarboxylic acid	772.1 $\pm$ 38.8	213.7 $\pm$ 14.6
(1S*, 2S*, 4S*, 5R*, 6S*) 2-Amino-4-benzoylamino-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid	511.5 $\pm$ 89	9.54 $\pm$ 0.73

To study the functional effect of compounds of the present invention in comparison to compounds of United

States Patent No. 5,958,960, forskolin stimulated c-AMP production in cells expressing recombinant mGlu2 and mGlu3 receptors was determined. As mGlu2 and mGlu3 receptors are negatively coupled to adenylyl cyclase, activity of an agonist decreases forskolin-stimulated accumulation of cyclic adenosine monophosphate (cAMP) and is reported as the concentration necessary to reduce C-AMP accumulation to 50% of the controls ( $EC_{50}$ , nM). The activity of an antagonist blocks the agonist induced decrease in forskolin-stimulated accumulation of cyclic adenosine monophosphate (cAMP) and is reported as the concentration necessary to inhibit the reduction of C-AMP accumulation to 50% of the controls ( $IC_{50}$ , nM). (Schoepp D. D., Johnson B. G., Wright R. A., Salhoff C. R., Mayne N. G., Wu S., Cockerham S. L., Burnett J. P., Belagaje R., Bleakman D., and Monn J., LY354740 is a potent and highly selective group II metabotropic glutamate receptor agonist in cells expressing human glutamate receptors. *Neuropharmacology* 36: 1-11 (1997)). As shown in table 2, compounds of the current invention have agonist effects on cells expressing mGlu2 and mGlu3 receptors where as compounds of United States Patent No. 5,958,960 have antagonist effects on cells expressing mGlu2 and mGlu3 receptors.

Table 2. Effects on forskolin-stimulated accumulation of cyclic adenosine monophosphate (cAMP) in cells expressing mGlu2 and mGlu3 receptors.		
Compounds of U.S. 5,958,960	IC <sub>50</sub> (nM) $\pm$ SEM	
	mGlu2	mGlu3
(1S*, 2S*, 4R*, 5R*, 6S*) 2-Amino-4-(3-methylureido)bicyclo[3.1.0]hexane-2,6-dicarboxylic acid	3,220 $\pm$ 1,310	6,270 $\pm$ 200
(1S*, 2S*, 4R*, 5R*, 6S*) 4-Acetylamino-2-aminobicyclo[3.1.0]hexane-2,6-dicarboxylic acid	4,300 $\pm$ 1,240	11,130 $\pm$ 1,080
1S*, 2S*, 4R*, 5R*, 6S*) 2-Amino-4-benzoylamino-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid	4,060 $\pm$ 1,400	9,520 $\pm$ 1,230
Compounds of present invention	EC <sub>50</sub> (nM) $\pm$ SEM	
	mGlu2	mGlu3
(1S*, 2S*, 4S*, 5R*, 6S*) 2-Amino-4-(3-methylureido)bicyclo[3.1.0]hexane-2,6-dicarboxylic acid	426.0 $\pm$ 41.82	41.11 $\pm$ 11.10
(1S*, 2S*, 4S*, 5R*, 6S*) 4-Acetylamino-2-aminobicyclo[3.1.0]hexane-2,6-dicarboxylic acid	1,350 $\pm$ 380	1,730 $\pm$ 320
(1S*, 2S*, 4S*, 5R*, 6S*) 2-Amino-4-benzoylamino-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid	60.68 $\pm$ 1.03	18.15 $\pm$ 2.87

Each of the compounds of the current invention possess a primary amino group at the C2-position of the bicyclic ring and two carboxylic acid groups, one at the C2-position and one at the C6-position. In general, it has been found that ester and/or amide derivatives of these functional groups are inactive in the receptor binding test. However, it is believed that these compounds are converted in vivo to the corresponding amino diacid and can therefore function as pro-drugs. It will be appreciated that the present invention provides the active amino diacid as well as any pro-drug forms that are capable of generating the active amino diacid in vivo.

The compounds of the present invention are preferably formulated prior to administration. Therefore, another aspect of the present invention is a pharmaceutical formulation comprising a compound of formula I in combination with one or more pharmaceutically-acceptable carriers, diluents, or excipients. The present pharmaceutical formulations are prepared by known procedures using well-known and readily available ingredients. In making the compositions of the present invention, the active ingredient will usually be mixed with a carrier, or diluted by a carrier, or enclosed within a carrier, and may be in the form of a capsule, sachet, paper, or other container. When the carrier serves as a diluent, it may be a solid, semi-solid, or liquid material which acts as a vehicle, excipient, or medium for the active ingredient. The compositions can be in the form of tablets, pills, powders, lozenges, sachets, cachets, elixirs, suspensions, emulsions, solutions, syrups, aerosols, ointments containing, for example, up to 10% by weight of active compound, soft and hard gelatin capsules, suppositories, sterile injectable solutions, dermal patch, subcutaneous implant, and sterile packaged powders.

Some examples of suitable carriers, excipients, and diluents include lactose, dextrose, sucrose, sorbitol,

mannitol, starches, gum, acacia, calcium phosphate, alginates, tragacanth, gelatin, calcium silicate, microcrystalline cellulose, polyvinylpyrrolidone, cellulose, water syrup, methyl cellulose, methyl and propyl hydroxybenzoates, talc, magnesium stearate, stearic acid, and mineral oil. The formulations can additionally include lubricating agents, wetting agents (surfactants), emulsifying and suspending agents, preserving agents, sweetening agents, or flavoring agents. Compositions of the invention may be formulated so as to provide quick, sustained, or delayed release of the active ingredient after administration to the patient by employing procedures well known in the art.

The compositions are preferably formulated in a unit dosage form, each dosage containing from about 5 mg to about 500 mg, more preferably about 25 mg to about 300 mg of the active ingredient. As used herein, the term "active ingredient" refers to a compound included within the scope of formula I.

The term "unit dosage form" refers to a physically discrete unit suitable as unitary dosages for human subjects and other mammals, each unit containing a predetermined quantity of active material calculated to produce the desired therapeutic effect, in association with a suitable pharmaceutical carrier, diluent, or excipient.

The following Examples further illustrate the compounds of the present invention and the methods for their synthesis. The Examples are not intended to be limiting to the scope of the invention in any respect, and should not be so construed. All experiments were run under a positive pressure of dry nitrogen or argon. All solvents and reagents were purchased from commercial sources and used as received, unless otherwise indicated. Dry tetrahydrofuran (THF) was obtained by distillation from sodium or sodium benzophenone ketyl prior to use. Proton nuclear magnetic resonance ( $^1\text{H}$  NMR) spectra were obtained on a GE QE-300

spectrometer at 300.15 MHz, a Bruker AM-500 spectrometer at 500 MHz, or a Bruker AC-200P spectrometer at 200 MHz. Free atom bombardment mass spectroscopy (FABMS) was performed on a VG ZAB-2SE instrument. Field desorption mass spectroscopy (FDMS) was performed using either a VG 70SE or a Varian MAT 731 instrument. Optical rotations were measured with a Perkin-Elmer 241 polarimeter. Chromatographic separation on a Waters Prep 500 LC was generally carried out using a linear gradient of the solvents indicated in the text. The reactions were generally monitored for completion using thin layer chromatography (TLC). Thin layer chromatography was performed using E. Merck Kieselgel 60 F<sub>254</sub> plates, 5 cm X 10 cm, 0.25 mm thickness. Spots were detected using a combination of UV and chemical detection (plates dipped in a ceric ammonium molybdate solution [75 g of ammonium molybdate and 4 g of cerium (IV) sulfate in 500 mL of 10% aqueous sulfuric acid] and then heated on a hot plate). Flash chromatography was performed as described by Still, et al. Still, Kahn, and Mitra, *J. Org. Chem.*, 43, 2923 (1978). Elemental analyses for carbon, hydrogen, and nitrogen were determined on a Control Equipment Corporation 440 Elemental Analyzer, or were performed by the Universidad Complutense Analytical Centre (Facultad de Farmacia, Madrid, Spain). Melting points were determined in open glass capillaries on a Gallenkamp hot air bath melting point apparatus or a Büchi melting point apparatus, and are uncorrected.

The abbreviations, symbols and terms used in the examples have the following meanings.

Ac = acetyl  
AllocCl = allyl chloroformate  
Anal. = elemental analysis  
Bn or Bzl = benzyl  
Bu = butyl  
BOC = butoxycarbonyl  
calcd = calculated  
D<sub>2</sub>O = deuterium oxide



DCC = dicyclohexylcarbodiimide  
DIBAL-H = diisobutyl aluminum hydride  
DMAP = dimethylaminopyridine  
DMF = dimethylformamide  
DMSO = dimethylsulfoxide  
EDCI = N-ethyl-N'-N'-dimethylaminopropyl  
carbodiimide  
Et = ethyl  
EtOAc = ethyl acetate  
EtOH = ethanol  
FAB = Fast Atom Bombardment (Mass Spectrascopy)  
FDMS = field desorption mass spectrum  
HOAt = 1-hydroxy-7-azabenzotriazole  
HOBt = 1-hydroxybenzotriazole  
HPLC = High Performance Liquid Chromatography  
HRMS = high resolution mass spectrum  
i-PrOH = isopropanol  
IR = Infrared Spectrum  
L = liter  
Me = methyl  
MeOH = methanol  
MPLC = Medium Pressure Liquid Chromatography  
Mp = melting point  
MTBE = t-butyl methyl ether  
NaHMDS = sodium hexamethyldisilylamide  
NBS = N-bromosuccinimide  
NMDBA = 1,3-dimethylbarbituric acid  
NMR = Nuclear Magnetic Resonance  
p-TsCl = para-toluenesulfonyl chloride  
p-TsOH = para-toulenesulfonic acid  
Ph = phenyl  
p.o. = oral administration  
i-Pr = isopropyl  
Rochelle's Salt = potassium sodium tartrate  
SM = starting material  
TBS = tert-butyldimethylsilyl


- 33 -

TEA = triethylamine  
Temp. = temperature  
TFA = trifluoroacetic acid  
THF = tetrahydrofuran  
TLC = thin layer chromatography  
t-BOC = tert-butoxycarbonyl

### Experimental Procedures

#### Preparation 1

(1*S*\*,2*S*\*,4*S*\*,5*R*\*,6*S*\*) Diethyl 2-(N-t-butyloxycarbonylamino)-4-aminobicyclo-[3.1.0]hexane-2,6-dicarboxylic acid



#### Method A

A. (1*S*\*,2*S*\*,4*S*\*,5*R*\*,6*R*\*) Diethyl 2-(N-t-butyloxy-carbonylamino)-4-(p-toluenesulfonyloxy)bicyclo[3.1.0]hexane-2,6-dicarboxylic acid.

p-Toluenesulfonyl chloride (21.3 g, 112 mmol) was added portionwise to a 0° C solution of (1*S*\*,2*S*\*,4*S*\*,5*R*\*,6*R*\*) diethyl 2-(N-t-butyloxycarbonylamino)-4-hydroxybicyclo-[3.1.0]hexane-2,6-dicarboxylic acid (20.0 g, 56 mmol—synthesis previously described in US 5,958,960 example 8a) in pyridine (80 mL). Upon complete addition the reaction mixture was allowed to warm to room temperature as it stirred over the weekend. The reaction mixture was diluted with EtOAc (2L), washed with cold aqueous NaHSO<sub>4</sub> (3X) then brine, dried over MgSO<sub>4</sub>, concentrated in vacuo and

purified by SiO<sub>2</sub> chromatography (HPLC: 10 % EtOAc/hexanes to 50 % EtOAc/hexanes) to yield 26.8 g (52.4 mmol, 94 %) of a white foam.

FDMS:  $M^+ + 1 = 512$ .

Anal. calcd. for C<sub>24</sub>H<sub>33</sub>NO<sub>5</sub>S: C, 56.35; H, 6.50; N, 2.74.

Found: C, 56.13; H, 6.37; N, 2.78.

B. (1S\*,2S\*,4R\*,5R\*,6R\*) Diethyl 2-(N-t-butyloxy-carbonylamino)-4-hydroxybicyclo-[3.1.0]hexane-2,6-dicarboxylic acid.

To a room temperature solution of the product from 1a (1.38 g, 2.7 mmol) in dimethyl sulfoxide (25 mL) was added potassium nitrite (2.30 g, 27 mmol) in one portion. The resulting reaction mixture was stirred at 50° C overnight. The reaction mixture was cooled to room temperature, diluted with EtOAc (500 mL), washed with water (3X) then brine, dried over MgSO<sub>4</sub>, concentrated in vacuo and purified by SiO<sub>2</sub> chromatography (PC-TLC: 10 % EtOAc/hexanes to 20 % EtOAc/hexanes) to yield 0.61 g (1.71 mmol, 63 %) of a white foam.

FDMS:  $M^+ + 1 = 358$ .

Anal. calcd. for C<sub>17</sub>H<sub>27</sub>NO<sub>7</sub>: C, 57.13; H, 7.61; N, 3.92.

Found: C, 57.06; H, 7.70; N, 3.81.

C. (1S\*,2S\*,4R\*,5R\*,6R\*) Diethyl 2-(N-t-butyloxy-carbonylamino)-4-(p-toluenesulfonyloxy)bicyclo[3.1.0]hexane-2,6-dicarboxylic acid.

p-Toluenesulfonyl chloride (3.8 g, 20 mmol) was added in one portion to a 0° C solution of the product from 1b

-35-

(3.6 g, 10 mmol) in pyridine (15 mL). Upon complete addition the reaction mixture was allowed to warm to room temperature as it stirred overnight. The reaction mixture was diluted with EtOAc (500 mL), washed with cold aqueous NaHSO<sub>4</sub> (3X) then brine, dried over MgSO<sub>4</sub>, concentrated in vacuo and purified by SiO<sub>2</sub> chromatography (HPLC: 10 % EtOAc/hexanes to 50 % EtOAc/hexanes) to yield 4.71 g (9.2 mmol, 92 %) of a white foam.

FDMS:  $M^+ + 1 = 512$ .

Anal. calcd. for C<sub>24</sub>H<sub>33</sub>NO<sub>9</sub>S: C, 56.35; H, 6.50; N, 2.74.

Found: C, 55.99; H, 6.15; N, 2.78.

D. (1S\*,2S\*,4S\*,5R\*,6S\*) Diethyl 2-(N-t-butyloxy-carbonylamino)-4-azidobicyclo-[3.1.0]hexane-2,6-dicarboxylic acid.

To a room temperature solution of the product from 1c (1.0 g, 2.0 mmol) in dimethyl sulfoxide (10 mL) was added sodium azide (0.39 g, 6.0 mmol) in one portion. The resulting reaction mixture was stirred at 45 °C overnight. The reaction mixture was cooled to room temperature, diluted with EtOAc (500 mL), washed with water (3X) then brine, dried over MgSO<sub>4</sub>, concentrated in vacuo and purified by SiO<sub>2</sub> chromatography (PC-TLC: 10 % EtOAc/hexanes to 20 % EtOAc/hexanes) to yield 0.76 g (2.0 mmol, 86 %) of a white foam.

FDMS:  $M^+ + 1 = 383$ .

Anal. calcd. for C<sub>17</sub>H<sub>26</sub>N<sub>4</sub>O<sub>6</sub>: C, 53.39; H, 6.85; N, 14.65.

Found: C, 53.39; H, 6.93; N, 14.36.

E. (1S\*,2S\*,4S\*,5R\*,6S\*) Diethyl 2-(N-t-butyloxy-carbonylamino)-4-aminobicyclo-[3.1.0]hexane-2,6-dicarboxylic acid.

To a room temperature solution of the product from 1d (0.65 g, 1.7 mmol) in anhydrous tetrahydrofuran (15 mL) was added triphenylphosphine (0.53 g, 2.0 mmol) in one portion. The resulting reaction mixture was stirred at room temperature overnight. Water (2 mL) was added to the reaction mixture and subsequently stirred overnight at room temperature. The reaction mixture was partitioned between saturated aqueous sodium bicarbonate and EtOAc. The product was extracted with EtOAc. All organics were combined, washed with water then brine, dried over  $K_2CO_3$ , concentrated in vacuo and purified by  $SiO_2$  chromatography (PC-TLC: 10 % EtOAc/hexanes to 90 % EtOAc/hexanes) to yield 0.56 g (1.57 mmol, 92 %) of a white foam.

FDMS:  $M^+ + 1 = 357$ .

Anal. calcd. for  $C_{17}H_{28}N_2O_6$ : C, 57.29; H, 7.92; N, 7.86.

Found: C, 57.00; H, 7.70; N, 7.65.

#### Method B

A. (1S\*,2S\*,4R\*,5R\*,6R\*) Diethyl 2-(N-t-butyloxy-carbonylamino)-4-bromobicyclo-[3.1.0]hexane-2,6-dicarboxylic acid.

Bromine was added dropwise to a room temperature solution of triphenylphosphine (0.40 g, 1.5 mmol) in  $CH_2Cl_2$  (25 mL) until a light yellow color persisted. Additional triphenylphosphine was added to the reaction mixture until the solution became colorless. The reaction mixture was chilled to 0 °C and a solution of (1S\*,2S\*,4S\*,5R\*,6R\*)

-37-

diethyl 2-(N-t-butyloxycarbonylamino)-4-hydroxy-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid (0.36 g, 1.0 mmol) in a 1:1 mixture of  $\text{CH}_2\text{Cl}_2$ /pyridine (20 mL total volume) was added dropwise to the reaction mixture. Upon complete addition the reaction mixture was allowed to warm to room temperature as it stirred overnight. The reaction mixture was diluted with  $\text{Et}_2\text{O}$  (500 mL), washed with cold aqueous  $\text{NaHSO}_4$  (3X) then brine, dried over  $\text{MgSO}_4$ , concentrated in vacuo and purified by  $\text{SiO}_2$  chromatography (PC-TLC: 10 %  $\text{EtOAc}$ /hexanes to 50 %  $\text{EtOAc}$ /hexanes) to yield 0.41 g (0.98 mmol, 97 %) of a white foam.

FDMS:  $\text{M}^+ + 18 (\text{NH}_4^+) = 439$ .

Anal. calcd. for  $\text{C}_{17}\text{H}_{26}\text{BrNO}_6$ : C, 48.58; H, 6.24; N, 3.33.

Found: C, 48.31; H, 6.04; N, 3.15.

B. (1S\*,2S\*,4S\*,5R\*,6S\*) Diethyl 2-(N-t-butyloxy-carbonylamino)-4-azidobicyclo[3.1.0]hexane-2,6-dicarboxylic acid.

To a room temperature solution of the product from 1f (13.2 g, 31.4 mmol) in dimethyl sulfoxide (150 mL) was added sodium azide (4.1 g, 62. mmol) in one portion. The resulting reaction mixture was stirred at 50°C overnight. The reaction mixture was cooled to room temperature, diluted with  $\text{EtOAc}$  (2 L), washed with water (3X) then brine, dried over  $\text{MgSO}_4$ , concentrated in vacuo and purified by  $\text{SiO}_2$  chromatography (HPLC: 10 %  $\text{EtOAc}$ /hexanes to 50 %  $\text{EtOAc}$ /hexanes) to yield 11.3 g (29.5 mmol, 94 %) of a white foam.

FDMS:  $\text{M}^+ + 1 = 383$ .

Anal. calcd. for  $\text{C}_{17}\text{H}_{26}\text{N}_4\text{O}_6$ : C, 53.39; H, 6.85; N, 14.65.

Found: C, 53.28; H, 6.79; N, 14.41.

C. (1S\*,2S\*,4S\*,5R\*,6S\*) Diethyl 2-(N-t-butyloxy-carbonylamino)-4-aminobicyclo-[3.1.0]hexane-2,6-dicarboxylic acid.

To a room temperature solution of the product from 1d (0.65 g, 1.7 mmol) in anhydrous tetrahydrofuran (15 mL) was added triphenylphosphine (0.53 g, 2.0 mmol) in one portion. The resulting reaction mixture was stirred at room temperature overnight. Water (2 mL) was added to the reaction mixture and subsequently stirred overnight at room temperature. The reaction mixture was partitioned between saturated aqueous sodium bicarbonate and EtOAc. The product was extracted with EtOAc. All organics were combined, washed with water then brine, dried over  $K_2CO_3$ , concentrated in vacuo and purified by  $SiO_2$  chromatography (PC-TLC: 10 % EtOAc/hexanes to 90 % EtOAc/hexanes) to yield 0.56 g (1.57 mmol, 92 %) of a white foam.


FDMS:  $M^+ + 1 = 357$ .

Anal. calcd. for  $C_{17}H_{28}N_2O_6$ : C, 57.29; H, 7.92; N, 7.86.

Found: C, 57.00; H, 7.70; N, 7.65.

#### Example 1

Synthesis of (1S\*,2S\*,4S\*,5R\*,6S\*) 2-amino-4-[3-(2-methoxyphenyl)ureido]bicyclo[3.1.0]hexane-2,6-dicarboxylic acid



A. (1S\*,2S\*,4S\*,5R\*,6S\*) Diethyl 2-(N-t-butyloxy-carbonylamino)-4-[3-(2-methoxy-phenyl)ureido]bicyclo[3.1.0]-hexane-2,6-dicarboxylic acid.

A room temperature solution of the product of Preparation 1 (0.25 g, 0.70 mmol) in  $\text{CH}_2\text{Cl}_2$  (10 mL) was treated in one portion with 2-methoxyphenyl isocyanate (0.13 g, 0.85 mmol). The resulting reaction mixture was allowed to stir at room temperature overnight. The reaction mixture was concentrated in vacuo and purified by  $\text{SiO}_2$  chromatography (PC-TLC: 10 % EtOAc/hexanes to 90 % EtOAc/hexanes) to yield 0.34 g (0.67 mmol, 96 %) of a white foam.

FDMS:  $M^+ + 1 = 506$ .

Anal. calcd. for  $\text{C}_{25}\text{H}_{35}\text{N}_3\text{O}_8 \cdot 0.1 \text{ eq. EtOAc}$ : C, 59.31; H, 7.02; N, 8.17.

Found: C, 58.97; H, 6.94; N, 8.06.

B. (1S\*,2S\*,4S\*,5R\*,6S\*) 2-amino-4-[3-(2-methoxyphenyl)-ureido]bicyclo-[3.1.0]hexane-2,6-dicarboxylic acid.

A 0 °C solution of the product of example 1a ( 0.31 g, 0.61 mmol) in EtOAc (35mL) was purged with anhydrous HCl gas until the solution was saturated. The resulting reaction mixture was stirred at 0 °C for 4 hours and concentrated to



dryness in vacuo to afford the HCl salt. The resulting solids were reconstituted in a 1:1 mixture of 1N NaOH:THF (10 mL total volume) and stirred at room temperature overnight. The reaction mixture was adjusted to pH = 7 with 1N HCl and concentrated to dryness. The solids were reconstituted in water, adjusted to pH = 12 with 1N NaOH and purified by anion exchange chromatography (Bio-Rad® AG1-X8 resin; acetate form converted to hydroxide form). The product was eluted from the column with 3N AcOH and concentrated to dryness in vacuo to afford crude product. Trituration in 2-propanol:water (5:1) followed by drying under vacuum at 80 °C afforded 0.20 g (0.57 mmol, 94 %) of the desired product as a white solid.

mp = >250 °C.


FDMS:  $M^+ + 1 = 350$ .

Anal. calcd. for  $C_{16}H_{19}N_3O_6 \cdot 1.0$  eq.  $H_2O$ : C, 52.31; H, 5.76; N, 11.44.

Found: C, 52.58; H, 5.52; N, 11.49.

#### Example 2

Synthesis of (1S\*,2S\*,4S\*,5R\*,6S\*) 2-amino-4-[3-(3-methoxyphenyl)ureido]bicyclo[3.1.0]hexane-2,6-dicarboxylic acid



-41-

A. (1S\*,2S\*,4S\*,5R\*,6S\*) Diethyl 2-(N-t-butyloxycarbonyl-amino)-4-[3-(3-methoxy-phenyl)ureido]bicyclo[3.1.0]hexane-2,6-dicarboxylic acid.

A room temperature solution of the product of Preparation 1 (0.25 g, 0.70 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (10 mL) was treated in one portion with 3-methoxyphenyl isocyanate (0.13 g, 0.85 mmol). The resulting reaction mixture was allowed to stir at room temperature overnight. The reaction mixture was concentrated in vacuo and purified by SiO<sub>2</sub> chromatography (PC-TLC: 10 % EtOAc/hexanes to 90 % EtOAc/hexanes) to yield 0.32 g (0.63 mmol, 90 %) of a white foam.

FDMS: M<sup>+</sup> + 1 = 506.

Anal. calcd. for C<sub>25</sub>H<sub>35</sub>N<sub>3</sub>O<sub>8</sub>·0.1 eq. CH<sub>2</sub>Cl<sub>2</sub>: C, 58.64; H, 6.90; N, 8.17.

Found: C, 58.36; H, 6.74; N, 8.06.

B. (1S\*,2S\*,4S\*,5R\*,6S\*) 2-amino-4-[3-(3-methoxyphenyl)-ureido]bicyclo-[3.1.0]hexane-2,6-dicarboxylic acid.

A 0 °C solution of the product of Part A ( 0.30 g, 0.60 mmol) in EtOAc (35mL) was purged with anhydrous HCl gas until the solution was saturated. The resulting reaction mixture was stirred at 0°C for 4 hours and concentrated to dryness in vacuo to afford the HCl salt. The resulting solids were reconstituted in a 1:1 mixture of 1N NaOH:THF (10 mL total volume) and stirred at room temperature overnight. The reaction mixture was adjusted to pH = 7 with 1N HCl and concentrated to dryness. The solids were reconstituted in water, adjusted to pH = 12 with 1N NaOH and purified by anion exchange chromatography (Bio-Rad® AG1-X8 resin; acetate form converted to hydroxide form). The

-42-

product was eluted from the column with 3N AcOH and concentrated to dryness in vacuo to afford crude product. Trituration in 2-propanol:water (5:1) followed by drying under vacuum at 80 °C afforded 0.16 g (0.46 mmol, 76 %) of the desired product as a white solid.

mp = >250 °C.


FDMS:  $M^+ + 1 = 350$ .

Anal. calcd. for  $C_{16}H_{19}N_3O_6 \cdot 0.3 \text{ eq. } H_2O \cdot 0.4 \text{ eq. } 2\text{-propanol}$ : C, 54.54; H, 6.07; N, 11.09.

Found: C, 54.39; H, 5.84; N, 10.82.

### Example 3

Synthesis of (1S\*,2S\*,4S\*,5R\*,6S\*) 2-amino-4-[3-(4-methoxyphenyl)ureido]bicyclo[3.1.0]hexane-2,6-dicarboxylic acid



A. (1S\*,2S\*,4S\*,5R\*,6S\*) Diethyl 2-(N-t-butyloxycarbonyl-amino)-4-[3-(4-methoxy-phenyl)ureido]bicyclo[3.1.0]hexane-2,6-dicarboxylic acid.

A room temperature solution of the product of preparation 1 (0.25 g, 0.70 mmol) in  $CH_2Cl_2$  (10 mL) was treated in one portion with 4-methoxyphenyl isocyanate (0.13 g, 0.85 mmol). The resulting reaction mixture was allowed to stir

at room temperature overnight. The reaction mixture was concentrated in vacuo and purified by SiO<sub>2</sub> chromatography (PC-TLC: 10 % EtOAc/hexanes to 90 % EtOAc/hexanes) to yield 0.30 g (0.60 mmol, 85 %) of a white foam.

FDMS:  $M^+ + 1 = 506$ .

Anal. calcd. for C<sub>25</sub>H<sub>35</sub>N<sub>3</sub>O<sub>8</sub>: C, 59.39; H, 6.98; N, 8.31.

Found: C, 58.36; H, 6.74; N, 8.06.

B. (1S\*,2S\*,4S\*,5R\*,6S\*) 2-amino-4-[3-(4-methoxyphenyl)-ureido]bicyclo[3.1.0]-hexane-2,6-dicarboxylic acid.

A 0 °C solution of the product of example 3a ( 0.30 g, 0.60 mmol) in EtOAc (35mL) was purged with anhydrous HCl gas until the solution was saturated. The resulting reaction mixture was stirred at 0°C for 4 hours and concentrated to dryness in vacuo to afford the HCl salt. The resulting solids were reconstituted in a 1:1 mixture of 1N NaOH:THF (15 mL total volume) and stirred at room temperature overnight. The reaction mixture was adjusted to pH = 7 with 1N HCl and concentrated to dryness. The solids were reconstituted in water, adjusted to pH = 12 with 1N NaOH and purified by anion exchange chromatography (Bio-Rad® AG1-X8 resin; acetate form converted to hydroxide form). The product was eluted from the column with 3N AcOH and concentrated to dryness in vacuo to afford crude product. Trituration in acetone:water (5:1) followed by drying under vacuum at 80 °C afforded 0.12 g (0.34 mmol, 62 %) of the desired product as a white solid.

mp = >250 °C.

FDMS:  $M^+ + 1 = 350$ .


-44-

Anal. calcd. for  $C_{16}H_{19}N_3O_6 \cdot 0.4 \text{ eq. } H_2O$ : C, 53.90; H, 5.60;  
N, 11.79.

Found: C, 53.76; H, 5.14; N, 11.43.

Example 4

Synthesis of (1S\*,2S\*,4S\*,5R\*,6S\*) 2-amino-4-[3-(2-methylphenyl)ureido]bicyclo[3.1.0]hexane-2,6-dicarboxylic acid



A. (1S\*,2S\*,4S\*,5R\*,6S\*) Diethyl 2-(N-t-butyloxycarbonyl-amino)-4-[3-(2-methyl-phenyl)ureido]bicyclo[3.1.0]hexane-2,6-dicarboxylic acid.

A room temperature solution of the product of preparation 1 (0.25 g, 0.70 mmol) in  $\text{CH}_2\text{Cl}_2$  (10 mL) was treated in one portion with 2-methylphenyl isocyanate (0.12 g, 0.85 mmol). The resulting reaction mixture was allowed to stir at room temperature overnight. The reaction mixture was concentrated in vacuo and purified by  $\text{SiO}_2$  chromatography (PC-TLC: 10 % EtOAc/hexanes to 90 % EtOAc/hexanes) to yield 0.32 g (0.65 mmol, 93 %) of a white foam.

FDMS:  $\text{M}^+ + 1 = 490$ .

Anal. calcd. for  $\text{C}_{25}\text{H}_{35}\text{N}_3\text{O}_7 \cdot 0.1 \text{ eq. } \text{CH}_2\text{Cl}_2$ : C, 60.53; H, 7.12; N, 8.44.

Found: C, 60.61; H, 7.01; N, 8.27.

B. (1S\*,2S\*,4S\*,5R\*,6S\*) 2-amino-4-[3-(2-methylphenyl)-ureido]bicyclo[3.1.0]hexane-2,6-dicarboxylic acid.

A 0°C solution of the product of example 4a ( 0.30 g, 0.60 mmol) in EtOAc (35mL) was purged with anhydrous HCl gas until the solution was saturated. The resulting reaction mixture was stirred at 0°C for 4 hours and concentrated to dryness in vacuo to afford the HCl salt. The resulting solids were reconstituted in a 1:1 mixture of 1N NaOH:THF (15 mL total volume) and stirred at room temperature overnight. The reaction mixture was adjusted to pH = 7 with 1N HCl and concentrated to dryness. The solids were reconstituted in water, adjusted to pH = 12 with 1N NaOH and purified by anion exchange chromatography (Bio-Rad® AG1-X8 resin; acetate form converted to hydroxide form). The product was eluted from the column with 3N AcOH and concentrated to dryness in vacuo to afford crude product. Trituration in 2-propanol:water (5:1) followed by drying under vacuum at 80 °C afforded 0.12 g (0.34 mmol, 62 %) of the desired product as a white solid.

mp = >250 °C.


FDMS:  $M^+ + 1 = 334$ .

Anal. calcd. for  $C_{16}H_{19}N_3O_5 \cdot 1.1 \text{ eq. } H_2O \cdot 0.3 \text{ eq. } AcOH$ : C, 53.75; H, 6.07; N, 11.19.

Found: C, 53.40; H, 5.70; N, 11.03.

Example 5

Synthesis of (1S\*,2S\*,4S\*,5R\*,6S\*) 2-amino-4-[3-(3-methylphenyl)ureido]bicyclo[3.1.0]hexane-2,6-dicarboxylic acid



A. (1S\*,2S\*,4S\*,5R\*,6S\*) Diethyl 2-(N-t-butyloxycarbonyl-amino)-4-[3-(3-methylphenyl)ureido]bicyclo[3.1.0]hexane-2,6-dicarboxylic acid.

A room temperature solution of the product of preparation 1 (0.25 g, 0.70 mmol) in  $\text{CH}_2\text{Cl}_2$  (10 mL) was treated in one portion with 3-methylphenyl isocyanate (0.12 g, 0.85 mmol). The resulting reaction mixture was allowed to stir at room temperature overnight. The reaction mixture was concentrated in vacuo and purified by  $\text{SiO}_2$  chromatography (PC-TLC: 10 % EtOAc/hexanes to 90 % EtOAc/hexanes) to yield 0.31 g (0.63 mmol, 90 %) of a white foam.

FDMS:  $\text{M}^+ + 1 = 490$ .

Anal. calcd. for  $\text{C}_{25}\text{H}_{35}\text{N}_3\text{O}_7 \cdot 0.1 \text{ eq. CH}_2\text{Cl}_2$ : C, 60.53; H, 7.12; N, 8.44.

Found: C, 60.74; H, 7.14; N, 8.37.

B. (1S\*,2S\*,4S\*,5R\*,6S\*) 2-amino-4-[3-(3-methylphenyl)-ureido]bicyclo[3.1.0]hexane-2,6-dicarboxylic acid.



A 0°C solution of the product of example 5a ( 0.30 g, 0.60 mmol) in EtOAc (35mL) was purged with anhydrous HCl gas until the solution was saturated. The resulting reaction mixture was stirred at 0°C for 4 hours and concentrated to dryness in vacuo to afford the HCl salt. The resulting solids were reconstituted in a 1:1 mixture of 1N NaOH:THF (15 mL total volume) and stirred at room temperature overnight. The reaction mixture was adjusted to pH = 7 with 1N HCl and concentrated to dryness. The solids were reconstituted in water, adjusted to pH = 12 with 1N NaOH and purified by anion exchange chromatography (Bio-Rad® AG1-X8 resin; acetate form converted to hydroxide form). The product was eluted from the column with 3N AcOH and concentrated to dryness in vacuo to afford crude product. Trituration in 2-propanol:water (5:1) followed by drying under vacuum at 80 °C afforded 0.12 g (0.34 mmol, 62 %) of the desired product as a white solid.

mp = >250 °C.


FDMS:  $M^+ + 1 = 334$ .

Anal. calcd. for  $C_{16}H_{19}N_3O_5 \cdot 0.6 \text{ eq. } H_2O$ : C, 55.84; H, 5.92; N, 12.21.

Found: C, 55.75; H, 5.84; N, 12.13.

Example 6

Synthesis of (1S\*,2S\*,4S\*,5R\*,6S\*) 2-amino-4-[3-(4-methylphenyl)ureido]bicyclo[3.1.0]hexane-2,6-dicarboxylic acid



A. (1S\*,2S\*,4S\*,5R\*,6S\*) Diethyl 2-(N-t-butyloxycarbonyl-amino)-4-[3-(4-methyl-phenyl)ureido]bicyclo[3.1.0]hexane-2,6-dicarboxylic acid.

A room temperature solution of the product of preparation 1 (0.25 g, 0.70 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (10 mL) was treated in one portion with 4-methylphenyl isocyanate (0.12 g, 0.85 mmol). The resulting reaction mixture was allowed to stir at room temperature overnight. The reaction mixture was concentrated in vacuo and purified by SiO<sub>2</sub> chromatography (PC-TLC: 10 % EtOAc/hexanes to 20 % EtOAc/hexanes) to yield 0.34 g (0.69 mmol, 99 %) of a white foam.

FDMS: M<sup>+</sup> + 1 = 490.

Anal. calcd. for C<sub>25</sub>H<sub>35</sub>N<sub>3</sub>O<sub>7</sub>·0.1 eq. CH<sub>2</sub>Cl<sub>2</sub>: C, 60.53; H, 7.12; N, 8.44.

Found: C, 60.71; H, 7.20; N, 8.44.

B. (1S\*,2S\*,4S\*,5R\*,6S\*) 2-amino-4-[3-(3-methylphenyl)-ureido]bicyclo[3.1.0]hexane-2,6-dicarboxylic acid.

A 0°C solution of the product of example 6a ( 0.30 g, 0.60 mmol) in EtOAc (35mL) was purged with anhydrous HCl gas until the solution was saturated. The resulting reaction mixture was stirred at 0°C for 4 hours and concentrated to dryness in vacuo to afford the HCl salt. The resulting solids were reconstituted in a 1:1 mixture of 1N NaOH:THF (15 mL total volume) and stirred at room temperature overnight. The reaction mixture was adjusted to pH = 7 with 1N HCl and concentrated to dryness. The solids were reconstituted in water, adjusted to pH = 12 with 1N NaOH and purified by anion exchange chromatography (Bio-Rad® AG1-X8 resin; acetate form converted to hydroxide form). The product was eluted from the column with 3N AcOH and concentrated to dryness in vacuo to afford crude product. Trituration in 2-propanol:water (5:1) followed by drying under vacuum at 80 °C afforded 0.12 g (0.36 mmol, 59 %) of the desired product as a white solid.

mp = charred >220°C.


FDMS:  $M^+ + 1 = 334$ .

Anal. calcd. for  $C_{16}H_{19}N_3O_5$ : C, 57.65; H, 5.74; N, 12.60.

Found: C, 60.55; H, 6.52; N, 12.26.

Example 7

Synthesis of (1S\*,2S\*,4S\*,5R\*,6S\*) 2-amino-4-[3-(2-chlorophenyl)ureido]bicyclo[3.1.0]hexane-2,6-dicarboxylic acid



A. (1S\*,2S\*,4S\*,5R\*,6S\*) Diethyl 2-(N-t-butyloxycarbonyl-amino)-4-[3-(2-chloro-phenyl)ureido]bicyclo[3.1.0]hexane-2,6-dicarboxylic acid.

A room temperature solution of the product of preparation 1 (0.25 g, 0.70 mmol) in  $\text{CH}_2\text{Cl}_2$  (10 mL) was treated in one portion with 2-chlorophenyl isocyanate (0.15 g, 1.0 mmol). The resulting reaction mixture was allowed to stir at room temperature overnight. The reaction mixture was concentrated in vacuo and purified by  $\text{SiO}_2$  chromatography (PC-TLC: 10 % EtOAc/hexanes to 50 % EtOAc/hexanes) to yield 0.35 g (0.69 mmol, 98 %) of a white foam.

FDMS:  $M^+ + 1 = 510$ .

Anal. calcd. for  $\text{C}_{24}\text{H}_{32}\text{ClN}_3\text{O}_7 \cdot 0.1 \text{ eq. CH}_2\text{Cl}_2$ : C, 55.83; H, 6.26; N, 8.10.

Found: C, 55.95; H, 6.21; N, 8.14.

B. (1S\*,2S\*,4S\*,5R\*,6S\*) 2-amino-4-[3-(2-chlorophenyl)-ureido]bicyclo-[3.1.0]hexane-2,6-dicarboxylic acid.

-52-

A 0°C solution of the product of example 7a (0.30 g, 0.59 mmol) in EtOAc (35mL) was purged with anhydrous HCl gas until the solution was saturated. The resulting reaction mixture was stirred at 0°C for 4 hours and concentrated to dryness in vacuo to afford the HCl salt. The resulting solids were reconstituted in a 1:1 mixture of 1N NaOH:THF (15 mL total volume) and stirred at room temperature overnight. The reaction mixture was adjusted to pH = 7 with 1N HCl and concentrated to dryness. The solids were reconstituted in water, adjusted to pH = 12 with 1N NaOH and purified by anion exchange chromatography (Bio-Rad® AG1-X8 resin; acetate form converted to hydroxide form). The product was eluted from the column with 3N AcOH and concentrated to dryness in vacuo to afford crude product. Trituration in 2-propanol:water (5:1) followed by drying under vacuum at 80 °C afforded 0.04 g (0.11 mmol, 19 %) of the desired product as a white solid.

mp = >250 °C.


FDMS: M - 1 = 352.

Anal. calcd. for  $C_{15}H_{16}ClN_3O_5 \cdot 1.1 \text{ eq. } H_2O$ : C, 48.23; H, 4.91; N, 11.25.

Found: C, 48.00; H, 4.39; N, 11.09.

Example 8

Synthesis of (1S\*,2S\*,4S\*,5R\*,6S\*) 2-amino-4-[3-(3-chlorophenyl)ureido]bicyclo[3.1.0]hexane-2,6-dicarboxylic acid



A. (1S\*,2S\*,4S\*,5R\*,6S\*) Diethyl 2-(N-t-butyloxycarbonyl-amino)-4-[3-(3-chloro-phenyl)ureido]bicyclo[3.1.0]hexane-2,6-dicarboxylic acid.

A room temperature solution of the product of preparation 1 (0.25 g, 0.70 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (10 mL) was treated in one portion with 3-chlorophenyl isocyanate (0.15 g, 1.0 mmol). The resulting reaction mixture was allowed to stir at room temperature overnight. The reaction mixture was concentrated in vacuo and purified by SiO<sub>2</sub> chromatography (PC-TLC: 10 % EtOAc/hexanes to 50 % EtOAc/hexanes) to yield 0.30 g (0.59 mmol, 84 %) of a white foam.

FDMS: M<sup>+</sup> + 1 = 510.

Anal. calcd. for C<sub>24</sub>H<sub>32</sub>ClN<sub>3</sub>O<sub>7</sub>·0.1 eq. CH<sub>2</sub>Cl<sub>2</sub>: C, 55.83; H, 6.26; N, 8.10.

Found: C, 55.63; H, 6.31; N, 8.02.

B. (1S\*,2S\*,4S\*,5R\*,6S\*) 2-amino-4-[3-(3chlorophenyl)-ureido]bicyclo[3.1.0]hexane-2,6-dicarboxylic acid.

-54-

A 0°C solution of the product of example 8a (0.25 g, 0.49 mmol) in EtOAc (35mL) was purged with anhydrous HCl gas until the solution was saturated. The resulting reaction mixture was stirred at 0°C for 4 hours and concentrated to dryness in vacuo to afford the HCl salt. The resulting solids were reconstituted in a 1:1 mixture of 1N NaOH:THF (15 mL total volume) and stirred at room temperature overnight. The reaction mixture was adjusted to pH = 7 with 1N HCl and concentrated to dryness. The solids were reconstituted in water, adjusted to pH = 12 with 1N NaOH and purified by anion exchange chromatography (Bio-Rad® AG1-X8 resin; acetate form converted to hydroxide form). The product was eluted from the column with 3N AcOH and concentrated to dryness in vacuo to afford crude product. Trituration in 2-propanol:water (5:1) followed by drying under vacuum at 80 °C afforded 0.06 g (0.17 mmol, 35 %) of the desired product as a white solid.

mp = >250 °C.


FDMS: M - 1 = 352.

Anal. calcd. for  $C_{15}H_{16}ClN_3O_5 \cdot 0.7 \text{ eq. } H_2O$ : C, 49.17; H, 4.79; N, 11.47.

Found: C, 48.96; H, 4.54; N, 11.23.

Example 9

Synthesis of (1S\*,2S\*,4S\*,5R\*,6S\*) 2-amino-4-[3-(4-chlorophenyl)ureido]bicyclo[3.1.0]hexane-2,6-dicarboxylic acid



A. (1S\*,2S\*,4S\*,5R\*,6S\*) Diethyl 2-(N-t-butyloxycarbonyl-amino)-4-[3-(4-chloro-phenyl)ureido]bicyclo[3.1.0]hexane-2,6-dicarboxylic acid.

A room temperature solution of the product of preparation 1 (0.25 g, 0.70 mmol) in  $\text{CH}_2\text{Cl}_2$  (10 mL) was treated in one portion with 4-chlorophenyl isocyanate (0.15 g, 1.0 mmol). The resulting reaction mixture was allowed to stir at room temperature overnight. The reaction mixture was concentrated in vacuo and purified by  $\text{SiO}_2$  chromatography (PC-TLC: 10 % EtOAc/hexanes to 50 % EtOAc/hexanes) to yield 0.29 g (0.57 mmol, 81 %) of a white foam.

FDMS:  $\text{M}^+ + 1 = 510$ .

Anal. calcd. for  $\text{C}_{24}\text{H}_{32}\text{ClN}_3\text{O}_7 \cdot 0.2 \text{ eq. } \text{CH}_2\text{Cl}_2$ : C, 55.16; H, 6.20; N, 7.97.

Found: C, 55.44; H, 6.19; N, 7.96.

B. (1S\*,2S\*,4S\*,5R\*,6S\*) 2-amino-4-[3-(4-chlorophenyl)-ureido]bicyclo[3.1.0]hexane-2,6-dicarboxylic acid.



A 0°C solution of the product of example 9a ( 0.25 g, 0.49 mmol) in EtOAc (35mL) was purged with anhydrous HCl gas until the solution was saturated. The resulting reaction mixture was stirred at 0°C for 4 hours and concentrated to dryness in vacuo to afford the HCl salt. The resulting solids were reconstituted in a 1:1 mixture of 1N NaOH:THF (15 mL total volume) and stirred at room temperature overnight. The reaction mixture was adjusted to pH = 7 with 1N HCl and concentrated to dryness. The solids were reconstituted in water, adjusted to pH = 12 with 1N NaOH and purified by anion exchange chromatography (Bio-Rad® AG1-X8 resin; acetate form converted to hydroxide form). The product was eluted from the column with 3N AcOH and concentrated to dryness in vacuo to afford crude product. Trituration in 2-propanol:water (5:1) followed by drying under vacuum at 80 °C afforded 0.05 g (0.15 mmol, 31 %) of the desired product as a white solid.

mp = >250 °C.


FDMS: M - 1 = 352.

Anal. calcd. for  $C_{15}H_{16}ClN_3O_5 \cdot 1.6 \text{ eq. } H_2O$ : C, 47.09; H, 5.06; N, 10.98.

Found: C, 46.86; H, 4.65; N, 10.72.

Example 10

Synthesis of (1S\*,2S\*,4S\*,5R\*,6S\*) 2-amino-4-[3-(2-fluorophenyl)ureido]-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid



A. (1S\*,2S\*,4S\*,5R\*,6S\*) Diethyl 2-(N-t-butyloxycarbonyl-amino)-4-[3-(2-fluoro-phenyl)ureido]bicyclo[3.1.0]hexane-2,6-dicarboxylic acid.

A room temperature solution of the product of preparation 1 (0.25 g, 0.70 mmol) in  $\text{CH}_2\text{Cl}_2$  (10 mL) was treated in one portion with 2-fluorophenyl isocyanate (0.12 g, 0.85 mmol). The resulting reaction mixture was allowed to stir at room temperature overnight. The reaction mixture was concentrated in vacuo and purified by  $\text{SiO}_2$  chromatography (PC-TLC: 10 % EtOAc/hexanes to 50 % EtOAc/hexanes) to yield 0.32 g (0.64 mmol, 93 %) of a white foam.

FDMS:  $\text{M}^+ + 1 = 494$ .

Anal. calcd. for  $\text{C}_{24}\text{H}_{32}\text{FN}_3\text{O}_7 \cdot 0.2 \text{ eq. CH}_2\text{Cl}_2$ : C, 56.93; H, 6.40; N, 8.23.

Found: C, 57.12; H, 6.16; N, 8.24.

B. (1S\*,2S\*,4S\*,5R\*,6S\*) 2-amino-4-[3-(2-fluorophenyl)-ureido]bicyclo[3.1.0]hexane-2,6-dicarboxylic acid.

A 0°C solution of the product of example 10a (0.25 g, 0.51 mmol) in EtOAc (35mL) was purged with anhydrous HCl gas until the solution was saturated. The resulting reaction mixture was stirred at 0°C for 4 hours and concentrated to dryness in vacuo to afford the HCl salt. The resulting solids were reconstituted in a 1:1 mixture of 1N NaOH:THF (15 mL total volume) and stirred at room temperature overnight. The reaction mixture was adjusted to pH = 7 with 1N HCl and concentrated to dryness. The solids were reconstituted in water, adjusted to pH = 12 with 1N NaOH and purified by anion exchange chromatography (Bio-Rad® AG1-X8 resin; acetate form converted to hydroxide form). The product was eluted from the column with 3N AcOH and concentrated to dryness in vacuo to afford crude product. Trituration in 2-propanol:water (5:1) followed by drying under vacuum at 80 °C afforded 0.10 g (0.30 mmol, 59 %) of the desired product as a white solid.

mp = >270 °C.


FDMS:  $M^+ + 1 = 338$ .

Anal. calcd. for  $C_{15}H_{16}FN_3O_5 \cdot 1.8 \text{ eq. } H_2O$ : C, 48.73; H, 5.34; N, 11.37.

Found: C, 48.56; H, 5.00; N, 11.59.

Example 11

Synthesis of (1S\*,2S\*,4S\*,5R\*,6S\*) 2-Amino-4-[3-(3-fluorophenyl)ureido]bicyclo[3.1.0]hexane-2,6-dicarboxylic acid



A. (1S\*,2S\*,4S\*,5R\*,6S\*) Diethyl 2-(N-t-butyloxycarbonyl-amino)-4-[3-(3-fluoro-phenyl)ureido]bicyclo[3.1.0]hexane-2,6-dicarboxylic acid.

A room temperature solution of the product of preparation 1 (0.25 g, 0.70 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (10 mL) was treated in one portion with 3-fluorophenyl isocyanate (0.12 g, 0.85 mmol). The resulting reaction mixture was allowed to stir at room temperature overnight. The reaction mixture was concentrated in vacuo and purified by SiO<sub>2</sub> chromatography (PC-TLC: 10 % EtOAc/hexanes to 50 % EtOAc/hexanes) to yield 0.34 g (0.69 mmol, 98 %) of a white foam.

FDMS: M<sup>+</sup> + 1 = 494.

Anal. calcd. for C<sub>24</sub>H<sub>32</sub>FN<sub>3</sub>O<sub>7</sub>·0.1 eq. CH<sub>2</sub>Cl<sub>2</sub>: C, 57.66; H, 6.47; N, 8.37.

Found: C, 57.40; H, 6.33; N, 8.18.

B. (1S\*,2S\*,4S\*,5R\*,6S\*) 2-amino-4-[3-(3-fluorophenyl)-ureido]bicyclo[3.1.0]hexane-2,6-dicarboxylic acid.

-60-

A 0°C solution of the product of example 11a (0.30 g, 0.61 mmol) in EtOAc (35mL) was purged with anhydrous HCl gas until the solution was saturated. The resulting reaction mixture was stirred at 0°C for 4 hours and concentrated to dryness in vacuo to afford the HCl salt. The resulting solids were reconstituted in a 1:1 mixture of 1N NaOH:THF (15 mL total volume) and stirred at room temperature overnight. The reaction mixture was adjusted to pH = 7 with 1N HCl and concentrated to dryness. The solids were reconstituted in water, adjusted to pH = 12 with 1N NaOH and purified by anion exchange chromatography (Bio-Rad® AG1-X8 resin; acetate form converted to hydroxide form). The product was eluted from the column with 3N AcOH and concentrated to dryness in vacuo to afford crude product. Trituration in 2-propanol:water (5:1) followed by drying under vacuum at 80 °C afforded 0.17 g (0.50 mmol, 83 %) of the desired product as a white solid.

mp = >250 °C.


FDMS:  $M^+ + 1 = 338$ .

Anal. calcd. for  $C_{15}H_{16}FN_3O_5 \cdot 1.5 \text{ eq. } H_2O$ : C, 49.45; H, 5.26; N, 11.53.

Found: C, 49.22; H, 5.18; N, 11.65.

Example 12

Synthesis of (1S\*,2S\*,4S\*,5R\*,6S\*) 2-amino-4-[3-(4-fluorophenyl)ureido]bicyclo[3.1.0]hexane-2,6-dicarboxylic acid



A. (1S\*,2S\*,4S\*,5R\*,6S\*) Diethyl 2-(N-t-butyloxycarbonyl-amino)-4-[3-(4-fluoro-phenyl)ureido]bicyclo[3.1.0]hexane-2,6-dicarboxylic acid.

A room temperature solution of the product of preparation 1 (0.25 g, 0.70 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (10 mL) was treated in one portion with 4-fluorophenyl isocyanate (0.12 g, 0.85 mmol). The resulting reaction mixture was allowed to stir at room temperature overnight. The reaction mixture was concentrated in vacuo and purified by SiO<sub>2</sub> chromatography (PC-TLC: 10 % EtOAc/hexanes to 50 % EtOAc/hexanes) to yield 0.35 g (0.70 mmol, 100 %) of a white foam.

FDMS: M<sup>+</sup> + 1 = 494.

Anal. calcd. for C<sub>24</sub>H<sub>32</sub>FN<sub>3</sub>O<sub>7</sub>·0.1 eq. CH<sub>2</sub>Cl<sub>2</sub>: C, 57.66; H, 6.47; N, 8.37.

Found: C, 57.55; H, 6.40; N, 8.24.

B. (1S\*,2S\*,4S\*,5R\*,6S\*) 2-amino-4-[3-(4-fluorophenyl)-ureido]bicyclo[3.1.0]hexane-2,6-dicarboxylic acid.

-62-

A 0 °C solution of the product of example 12a (0.30 g, 0.61 mmol) in EtOAc (35mL) was purged with anhydrous HCl gas until the solution was saturated. The resulting reaction mixture was stirred at 0 °C for 4 hours and concentrated to dryness in vacuo to afford the HCl salt. The resulting solids were reconstituted in a 1:1 mixture of 1N NaOH:THF (15 mL total volume) and stirred at room temperature overnight. The reaction mixture was adjusted to pH = 7 with 1N HCl and concentrated to dryness. The solids were reconstituted in water, adjusted to pH = 12 with 1N NaOH and purified by anion exchange chromatography (Bio-Rad® AG1-X8 resin; acetate form converted to hydroxide form). The product was eluted from the column with 3N AcOH and concentrated to dryness in vacuo to afford crude product. Trituration in 2-propanol:water (5:1) followed by drying under vacuum at 80 °C afforded 0.17 g (0.50 mmol, 83 %) of the desired product as a white solid.

mp = charred >220°C.


FDMS:  $M^+ + 1 = 338$ .

Anal. calcd. for  $C_{15}H_{16}FN_3O_5 \cdot 1.5 \text{ eq. } H_2O$ : C, 49.45; H, 5.26; N, 11.53.

Found: C, 49.08; H, 5.23; N, 11.61.

Example 13

Synthesis of (1S\*,2S\*,4S\*,5R\*,6S\*) 2-amino-4-[3-(2,6-dichlorophenyl)ureido]bicyclo[3.1.0]hexane-2,6-dicarboxylic acid



A. (1S\*,2S\*,4S\*,5R\*,6S\*) Diethyl 2-(N-t-butyloxycarbonyl-amino)-4-[3-(2,6-dichloro-phenyl)ureido]-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid.

A room temperature solution of the product of preparation 1 (0.25 g, 0.70 mmol) in  $\text{CH}_2\text{Cl}_2$  (10 mL) was treated in one portion with 2,6-dichlorophenyl isocyanate (0.16 g, 0.85 mmol). The resulting reaction mixture was allowed to stir at room temperature overnight. The reaction mixture was concentrated in vacuo and purified by  $\text{SiO}_2$  chromatography (PC-TLC: 10 % EtOAc/hexanes to 50 % EtOAc/hexanes) to yield 0.35 g (0.64 mmol, 92 %) of a white foam.

FDMS:  $M - 1 = 542$ .

Anal. calcd. for  $\text{C}_{24}\text{H}_{31}\text{Cl}_2\text{N}_3\text{O}_7 \cdot 0.1 \text{ eq. } \text{CH}_2\text{Cl}_2$ : C, 52.35; H, 5.69; N, 7.60.

Found: C, 52.34; H, 5.55; N, 7.46.



B. (1S\*,2S\*,4S\*,5R\*,6S\*) 2-amino-4-[3-(2,6-dichlorophenyl)-ureido]-bicyclo-[3.1.0]hexane-2,6-dicarboxylic acid.

A 0°C solution of the product of example 13a (0.33 g, 0.61 mmol) in EtOAc (35mL) was purged with anhydrous HCl gas until the solution was saturated. The resulting reaction mixture was stirred at 0°C for 4 hours and concentrated to dryness in vacuo to afford the HCl salt. The resulting solids were reconstituted in a 1:1 mixture of 1N NaOH:THF (15 mL total volume) and stirred at room temperature overnight. The reaction mixture was adjusted to pH = 7 with 1N HCl and concentrated to dryness. The solids were reconstituted in water, adjusted to pH = 12 with 1N NaOH and purified by anion exchange chromatography (Bio-Rad® AG1-X8 resin; acetate form converted to hydroxide form). The product was eluted from the column with 3N AcOH and concentrated to dryness in vacuo to afford crude product. Trituration in 2-propanol:water (5:1) followed by drying under vacuum at 80 °C afforded 0.54 g (0.54 mmol, 89 %) of the desired product as a white solid.

mp = >250 °C.


FDMS:  $M^+ + 1 = 388$ .

Anal. calcd. for  $C_{15}H_{15}Cl_2N_3O_5 \cdot 1.3 \text{ eq. } H_2O$ : C, 43.77; H, 4.31; N, 10.21.

Found: C, 43.55; H, 4.07; N, 10.04.

Example 14

Synthesis of (1S\*,2S\*,4S\*,5R\*,6S\*) 2-amino-4-[3-(2,4-dichlorophenyl)ureido]bicyclo[3.1.0]hexane-2,6-dicarboxylic acid



A. (1S\*,2S\*,4S\*,5R\*,6S\*) Diethyl 2-(N-t-butyloxycarbonyl-amino)-4-[3-(2,4-dichloro-phenyl)ureido]bicyclo[3.1.0]-hexane-2,6-dicarboxylic acid.

A room temperature solution of the product of preparation 1 (0.25 g, 0.70 mmol) in  $\text{CH}_2\text{Cl}_2$  (10 mL) was treated in one portion with 2,4-dichlorophenyl isocyanate (0.16 g, 0.85 mmol). The resulting reaction mixture was allowed to stir at room temperature overnight. The reaction mixture was concentrated in vacuo and purified by  $\text{SiO}_2$  chromatography (PC-TLC: 10 % EtOAc/hexanes to 50 % EtOAc/hexanes) to yield 0.36 g (0.66 mmol, 94 %) of a white foam.

FDMS:  $M - 1 = 542$ .

Anal. calcd. for  $\text{C}_{24}\text{H}_{31}\text{Cl}_2\text{N}_3\text{O}_7 \cdot 0.1 \text{ eq. } \text{CH}_2\text{Cl}_2$ : C, 52.35; H, 5.69; N, 7.60.

Found: C, 52.73; H, 5.76; N, 7.48.

B. (1S\*,2S\*,4S\*,5R\*,6S\*) 2-amino-4-[3-(2,4-dichlorophenyl)ureido]bicyclo-[3.1.0]hexane-2,4-dicarboxylic acid.

A 0°C solution of the product of example 14a (0.34 g, 0.62 mmol) in EtOAc (35mL) was purged with anhydrous HCl gas until the solution was saturated. The resulting reaction mixture was stirred at 0°C for 4 hours and concentrated to dryness in vacuo to afford the HCl salt. The resulting solids were reconstituted in a 1:1 mixture of 1N NaOH:THF (15 mL total volume) and stirred at room temperature overnight. The reaction mixture was adjusted to pH = 7 with 1N HCl and concentrated to dryness. The solids were reconstituted in water, adjusted to pH = 12 with 1N NaOH and purified by anion exchange chromatography (Bio-Rad® AG1-X8 resin; acetate form converted to hydroxide form). The product was eluted from the column with 3N AcOH and concentrated to dryness in vacuo to afford crude product. Trituration in 2-propanol:water (5:1) followed by drying under vacuum at 80 °C afforded 0.13 g (0.33 mmol, 53 %) of the desired product as a white solid.

mp = >250 °C.


FDMS: M - 1 = 388.

Anal. calcd. for C<sub>15</sub>H<sub>15</sub>Cl<sub>2</sub>N<sub>3</sub>O<sub>5</sub>: C, 46.41; H, 3.89; N, 10.82.

Found: C, 43.88; H, 3.87; N, 9.80.

Example 15

Synthesis of (1S\*,2S\*,4S\*,5R\*,6S\*) 2-amino-4-[3-(3,5-dichlorophenyl)ureido]bicyclo[3.1.0]hexane-2,6-dicarboxylic acid



A. (1S\*,2S\*,4S\*,5R\*,6S\*) Diethyl 2-(N-t-butyloxycarbonyl-amino)-4-[3-(3,5-dichloro-phenyl)ureido]bicyclo[3.1.0]-hexane-2,6-dicarboxylic acid.

A room temperature solution of the product of preparation 1 (0.25 g, 0.70 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (10 mL) was treated in one portion with 3,5-dichlorophenyl isocyanate (0.17 g, 0.88 mmol). The resulting reaction mixture was allowed to stir at room temperature overnight. The reaction mixture was concentrated in vacuo and purified by SiO<sub>2</sub> chromatography (PC-TLC: 10 % EtOAc/hexanes to 50 % EtOAc/hexanes) to yield 0.38 g (0.70 mmol, 99 %) of a white foam.

FDMS: M - 1 = 542.

Anal. calcd. for C<sub>24</sub>H<sub>31</sub>Cl<sub>2</sub>N<sub>3</sub>O<sub>7</sub>·0.1 eq. CH<sub>2</sub>Cl<sub>2</sub>: C, 52.35; H, 5.69; N, 7.60.

Found: C, 52.43; H, 5.69; N, 7.45.

B. (1S\*,2S\*,4S\*,5R\*,6S\*) 2-amino-4-[3-(3,5-dichlorophenyl)ureido]bicyclo-[3.1.0]hexane-3,5-dicarboxylic acid.

A 0°C solution of the product of example 15a (0.35 g, 0.64 mmol) in EtOAc (35mL) was purged with anhydrous HCl gas until the solution was saturated. The resulting reaction mixture was stirred at 0°C for 4 hours and concentrated to dryness in vacuo to afford the HCl salt. The resulting solids were reconstituted in a 1:1 mixture of 1N NaOH:THF (15 mL total volume) and stirred at room temperature overnight. The reaction mixture was adjusted to pH = 7 with 1N HCl and concentrated to dryness. The solids were reconstituted in water, adjusted to pH = 12 with 1N NaOH and purified by anion exchange chromatography (Bio-Rad® AG1-X8 resin; acetate form converted to hydroxide form). The product was eluted from the column with 3N AcOH and concentrated to dryness in vacuo to afford crude product. Trituration in methanol followed by drying under vacuum at 80 °C afforded 0.15 g (0.39 mmol, 61 %) of the desired product as a white solid.

mp = >250 °C.


FDMS: M - 2 = 386.

Anal. calcd. for C<sub>15</sub>H<sub>15</sub>Cl<sub>2</sub>N<sub>3</sub>O<sub>5</sub>·0.5 eq. H<sub>2</sub>O: C, 45.36; H, 4.06; N, 10.58.

Found: C, 45.11; H, 3.69; N, 10.88.

Example 16

Synthesis of (1S\*,2S\*,4S\*,5R\*,6S\*) 2-amino-4-[3-(3,4-dichlorophenyl)ureido]bicyclo[3.1.0]hexane-2,6-dicarboxylic acid



A. (1S\*,2S\*,4S\*,5R\*,6S\*) Diethyl 2-(N-t-butyloxycarbonyl-amino)-4-[3-(3,4-dichloro-phenyl)ureido]bicyclo[3.1.0]-hexane-2,6-dicarboxylic acid.

A room temperature solution of the product of preparation 1 (0.25 g, 0.70 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (10 mL) was treated in one portion with 3,4-dichlorophenyl isocyanate (0.17 g, 0.88 mmol). The resulting reaction mixture was allowed to stir at room temperature overnight. The reaction mixture was concentrated in vacuo and purified by SiO<sub>2</sub> chromatography (PC-TLC: 10 % EtOAc/hexanes to 50 % EtOAc/hexanes) to yield 0.36 g (0.66 mmol, 94 %) of a white foam.

FDMS: M - 1 = 542.

Anal. calcd. for C<sub>24</sub>H<sub>31</sub>Cl<sub>2</sub>N<sub>3</sub>O<sub>7</sub>·0.1 eq. CH<sub>2</sub>Cl<sub>2</sub>: C, 52.35; H, 5.69; N, 7.60.

Found: C, 52.26; H, 5.59; N, 7.34.

A 0 °C solution of the product of example 16a (0.33 g, 0.61 mmol) in EtOAc (35mL) was purged with anhydrous HCl gas until the solution was saturated. The resulting reaction mixture was stirred at 0 °C for 4 hours and concentrated to dryness in vacuo to afford the HCl salt. The resulting solids were reconstituted in a 1:1 mixture of 1N NaOH:THF (15 mL total volume) and stirred at room temperature overnight. The reaction mixture was adjusted to pH = 7 with 1N HCl and concentrated to dryness. The solids were reconstituted in water, adjusted to pH = 12 with 1N NaOH and purified by anion exchange chromatography (Bio-Rad® AG1-X8 resin; acetate form converted to hydroxide form). The product was eluted from the column with 3N AcOH and concentrated to dryness in vacuo to afford crude product. Trituration in methanol followed by drying under vacuum at 80 °C afforded 0.12 g (0.31 mmol, 51 %) of the desired product as a white solid.

mp = >250 °C.


FDMS: M - 2 = 386.

Anal. calcd. for  $C_{15}H_{15}Cl_2N_3O_5 \cdot 0.2 \text{ eq. } H_2O$ : C, 45.98; H, 3.96; N, 10.73.

Found: C, 45.73; H, 3.87; N, 10.42.

Example 17

Synthesis of (1S\*,2S\*,4S\*,5R\*,6S\*) 2-amino-4-[3-(2-naphthyl)ureido]bicyclo[3.1.0]hexane-2,6-dicarboxylic acid



A. (1S\*,2S\*,4S\*,5R\*,6S\*) Diethyl 2-(N-t-butyloxycarbonyl-amino)-4-[3-(2-naphthyl)-ureido]bicyclo[3.1.0]hexane-2,6-dicarboxylic acid.

A room temperature solution of the product of preparation 1 (0.25 g, 0.70 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (10 mL) was treated in one portion with 2-naphthyl isocyanate (0.15 g, 0.88 mmol). The resulting reaction mixture was allowed to stir at room temperature overnight. The reaction mixture was concentrated in vacuo and purified by SiO<sub>2</sub> chromatography (PC-TLC: 10 % EtOAc/hexanes to 50 % EtOAc/hexanes) to yield 0.36 g (0.68 mmol, 98 %) of a white foam.

FDMS: M<sup>+</sup> + 35 (Cl) = 561.

B. (1S\*,2S\*,4S\*,5R\*,6S\*) 2-amino-4-[3-(2-naphthyl)ureido]-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid.

A 0 °C solution of the product of example 18a (0.35 g, 0.67 mmol) in EtOAc (35mL) was purged with anhydrous HCl gas until the solution was saturated. The resulting reaction



-72-

mixture was stirred at 0 °C for 4 hours and concentrated to dryness in vacuo to afford the HCl salt. The resulting solids were reconstituted in a 1:1 mixture of 1N NaOH:THF (15 mL total volume) and stirred at room temperature overnight. The reaction mixture was adjusted to pH = 7 with 1N HCl and concentrated to dryness. The solids were reconstituted in water, adjusted to pH = 12 with 1N NaOH and purified by anion exchange chromatography (Bio-Rad® AG1-X8 resin; acetate form converted to hydroxide form). The product was eluted from the column with 3N AcOH and concentrated to dryness in vacuo to afford crude product. Trituration in acetone followed by drying under vacuum at 80 °C afforded 0.11 g (0.30 mmol, 43 %) of the desired product as a white solid.

mp = charred >250 °C.


FDMS:  $M^+ + 1 = 370$ .

Anal. calcd. for  $C_{19}H_{19}N_3O_5 \cdot 0.6$  eq. AcOH: C, 59.85; H, 5.32; N, 10.37.

Found: C, 59.74; H, 5.34; N, 10.54.

Example 18

Synthesis of (1S\*,2S\*,4S\*,5R\*,6S\*) 2-amino-4-[3-(1-naphthyl)ureido]bicyclo[3.1.0]hexane-2,6-dicarboxylic acid



A. (1S\*,2S\*,4S\*,5R\*,6S\*) Diethyl 2-(N-t-butyloxycarbonyl-amino)-4-[3-(1-naphthyl)-ureido]bicyclo[3.1.0]hexane-2,6-dicarboxylic acid.

A room temperature solution of the product of preparation 1 (0.25 g, 0.70 mmol) in  $\text{CH}_2\text{Cl}_2$  (10 mL) was treated in one portion with 1-naphthyl isocyanate (0.15 g, 0.88 mmol). The resulting reaction mixture was allowed to stir at room temperature overnight. The reaction mixture was filtered to remove insolubles, concentrated in vacuo and purified by  $\text{SiO}_2$  chromatography (PC-TLC: 10 % EtOAc/hexanes to 50 % EtOAc/hexanes) to yield 0.36 g (0.68 mmol, 98 %) of a white foam.

FDMS:  $\text{M}^+ + 1 = 526$ .

Anal. calcd. for  $\text{C}_{28}\text{H}_{35}\text{N}_3\text{O}_7$ : C, 63.99; H, 6.71; N, 7.99.

Found: C, 63.75; H, 6.72; N, 7.93.

B. (1S\*,2S\*,4S\*,5R\*,6S\*) 2-amino-4-[3-(1-naphthyl)ureido]-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid..

A 0 °C solution of the product of example 19a (0.31 g, 0.60 mmol) in EtOAc (35mL) was purged with anhydrous HCl gas until the solution was saturated. The resulting reaction mixture was stirred at 0 °C for 4 hours and concentrated to dryness in vacuo to afford the HCl salt. The resulting solids were reconstituted in a 1:1 mixture of 1N NaOH:THF (15 mL total volume) and stirred at room temperature overnight. The reaction mixture was adjusted to pH = 7 with 1N HCl and concentrated to dryness. The solids were reconstituted in water, adjusted to pH = 12 with 1N NaOH and purified by anion exchange chromatography (Bio-Rad® AG1-X8 resin; acetate form converted to hydroxide form). The product was eluted from the column with 3N AcOH and concentrated to dryness in vacuo to afford crude product. Trituration in 2-propanol/H<sub>2</sub>O (9:1) followed by drying under vacuum at 80 °C afforded 0.12 g (0.32 mmol, 45 %) of the desired product as a white solid.

mp = >250 °C.


FDMS:  $M^+ + 1 = 370$ .

Anal. calcd. for C<sub>19</sub>H<sub>19</sub>N<sub>3</sub>O<sub>5</sub>·1.0 eq. AcOH·0.3 eq. H<sub>2</sub>O: C, 58.00; H, 5.47; N, 9.66.

Found: C, 57.81; H, 5.09; N, 9.95.

Example 19

Synthesis of (1S\*, 2S\*, 4S\*, 5R\*, 6S\*) 2-Amino-4-(3-phenyl-ureido)-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid




The title compound was prepared utilizing the two step procedure in Example 1. In step (a) the product of Preparation 1 (0.34 g, 0.95 mmol) and phenyl isocyanate (0.12g, 1.0 mmol) were reacted to afford (1S\*, 2S\*, 4S\*, 5R\*, 6S\*)-2-tert-butoxycarbonylamino-4-(3-phenyl-ureido)-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid diethyl ester in 88.2% yield. In step (b) the intermediate urea (0.4 g, 0.84 mmol) was converted to the title compound and isolated as a solid in a 69% yield (0.186 g, 0.58 mmol) by cation exchange chromatography (Dowex 50X8-100: 10% Pyridine/H<sub>2</sub>O).

m.p. > 250°C.

FDMS: M<sup>+</sup>+1 = 320.

Example 20

Synthesis of (1S\*, 2S\*, 4S\*, 5R\*, 6S\*) 2-Amino-4-(3-methyl-ureido)-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid




The title compound was prepared utilizing the two step procedure in Example 1. In step (a) the product of Preparation 1 (0.25 g, 0.7 mmol) and methyl isocyanate (0.044 g, 0.77 mmol) were reacted to afford (1S\*, 2S\*, 4S\*, 5R\*, 6S\*)-2-tert-butoxycarbonylamino-4-(3-methyl-ureido)-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid diethyl ester in 73.7% yield. In step (b) the intermediate urea (0.2 g, 0.45 mmol) was converted to the title compound and isolated as a solid in a 95% yield (0.11 g, 0.43 mmol) by cation exchange chromatography (Dowex 50X8-100: 10% Pyridine/H<sub>2</sub>O).

m.p. > 250°C.

FDMS: M<sup>+</sup>+1 = 320.

Example 21

Synthesis of (1S\*,2S\*,4S\*,5R\*,6S\*) 2-Amino-4-(3-phenylthioureido)-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid



A. (1S\*,2S\*,4S\*,5R\*,6S\*) Diethyl 2-(N-t-butyloxycarbonyl-amino)-4-[3-phenylthioureido]bicyclo[3.1.0]hexane-2,6-dicarboxylic acid.

A room temperature solution of the product of preparation 1 (0.25 g, 0.70 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (10 mL) was treated in one portion with phenyl isothiocyanate (0.14 g, 1.0 mmol). The resulting reaction mixture was allowed to stir at room temperature overnight. The reaction mixture was concentrated in vacuo and purified by SiO<sub>2</sub> chromatography (PC-TLC: 10 % EtOAc/hexanes to 50 % EtOAc/hexanes) to yield 0.28 g (0.57 mmol, 81 %) of a white foam.

FDMS: M - 1 = 490.

Anal. calcd. for C<sub>24</sub>H<sub>33</sub>N<sub>3</sub>O<sub>6</sub>S\*0.1 eq. CH<sub>2</sub>Cl<sub>2</sub>: C, 57.88; H, 6.69; N, 8.40.

Found: C, 57.93; H, 6.52; N, 8.34.

B. (1S\*,2S\*,4S\*,5R\*,6S\*) 2-amino-4-[3-(1-naphthyl)ureido]-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid.

-78-

A 0 °C solution of the product of example 19a (0.31 g, 0.60 mmol) in EtOAc (35mL) was purged with anhydrous HCl gas until the solution was saturated. The resulting reaction mixture was stirred at 0 °C for 4 hours and concentrated to dryness in vacuo to afford the HCl salt. The resulting solids were reconstituted in a 1:1 mixture of 1N NaOH:THF (15 mL total volume) and stirred at room temperature overnight. The reaction mixture was adjusted to pH = 7 with 1N HCl and concentrated to dryness. The solids were reconstituted in water, adjusted to pH = 2 and the resulting precipitate collected via vacuum filtration to yield the title compound as the HCl salt. Subsequent drying under vacuum at 80 °C afforded 0.14 g (0.38 mmol, 71 %) of the desired product as a white solid.

mp = >275 °C.


FDMS:  $M^+ + 1 = 336$  (free base).

Anal. calcd. for  $C_{15}H_{17}N_3O_4S \cdot 1.0$  eq. HCl: C, 48.45; H, 4.88; N, 11.30.

Found: C, 48.58; H, 4.52; N, 11.17.

Example 22

Synthesis of (1S\*, 2S\*, 4S\*, 5R\*, 6S\*) 2-Amino-4-[(thiophene-3-carbonyl)-amino]-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid



A. (1S\*, 2S\*, 4S\*, 5R\*, 6S\*) 2-tert-Butoxycarbonylamino-4-[(thiophene-3-carbonyl)-amino]-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid diethyl ester.

3-Thiophenecarboxylic acid was dissolved in 3 ml of anhydrous DMF and cooled to 0°C under a nitrogen atmosphere. 1,3-Dicyclohexylcarbodiimide (0.133 g, 1.05 mmol) and hydroxybenztriazole (0.19 g, 1.4 mmol) were added sequentially to the acid. The reaction was allowed to warm to ambient temperature for 15 minutes, then cooled to 0°C again. The product of Preparation 1 (0.25 g, 0.7 mmol) in 2 ml of anhydrous DMF was added to the cooled reaction mixture above and allowed to stir over night. Partitioned the reaction between EtOAc and H<sub>2</sub>O, washed with brine and the organics dried over MgSO<sub>4</sub>. The isolated crude material was triturated with EtOAc and the solids were filtered. The product was purified using PT-TLC (10% EtOAc/hexanes to 50% EtOAc/hexanes) to give 100% yield (0.33 g, 0.7 mmol).

FDMS: M<sup>+</sup>-1 = 465.



B. (1S\*, 2S\*, 4S\*, 5R\*, 6S\*) 2-Amino-4-[(thiophene-3-carbonyl)-amino]-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid.

A solution of the product from step (a) (0.34 g, 0.73 mmol), in EtOAc was chilled to 0°C. The reaction solution was saturated with HCl (g) and allowed to react until the Boc group was cleaved as indicated by TLC. The reaction was concentrated under vacuum and rediluted in THF (10 ml). 1N NaOH (10 ml) was added and the mixture was allowed to stir overnight (pH = 14). The reaction mixture was titrated with 1N HCl to a pH = ~ 7 and organics were removed under vacuum. The title compound (0.118 g, 0.38 mmol) was isolated in 52.1% yield by isoelectric precipitation at pH = ~3-4.

m.p. > 250°C.


FDMS:  $M^+ + 1 = 311$ .

Anal. calcd. For  $C_{13}H_{14}N_2O_5S \cdot 0.2 H_2O$ : C, 49.74; H, 4.62; N, 8.92.

Found: C, 49.70; H, 4.59; N, 8.90.

Example 23

Synthesis of (1S\*, 2S\*, 4S\*, 5R\*, 6S\*) 4-Acetylamino-2-amino-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid



The title compound was prepared utilizing the two step procedure in Example 22. In step (a) the product of Preparation 1 (0.2 g, 0.56 mmol) and acetyl chloride (0.053 g, 0.67 mmol) were reacted to afford (1S\*, 2S\*, 4S\*, 5R\*, 6S\*)-4-acetylamino-2-tert-butoxycarbonylamino-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid diethyl ester in 99% yield. In step (b) the intermediate amide (0.19 g, 0.48 mmol) was converted to the title compound in a 61.9% yield (0.072 g, 0.3 mmol).

m.p. = 223-224°C.


FDMS:  $M^+ - 1 = 241$ .

Anal. calcd. For  $C_{10}H_{14}N_2O_5 \cdot 1.5 H_2O$ : C, 44.61; H, 6.36; N, 10.40.

Found: C, 44.67; H, 6.28; N, 10.80.

Example 24

Synthesis of (1S\*, 2S\*, 4S\*, 5R\*, 6S\*) 2-Amino-4-benzoylamino-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid




The title compound was prepared utilizing the two step procedure in Example 22. In step (a) the product of Preparation 1 (0.32 g, 0.8 mmol) and benzoyl chloride (0.15 g, 1.0 mmol) were reacted to afford (1S\*, 2S\*, 4S\*, 5R\*, 6S\*)-4-benzoylamino-2-tert-butoxycarbonylamino-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid diethyl ester in 99% yield. In step (b) the intermediate amide (0.39 g, 0.85 mmol) was converted to the title compound in a 46.6% yield (0.12 g, 0.39 mmol).

m.p. = 238-240°C.

FDMS: M<sup>+</sup>+1 = 305.

Example 25

Synthesis of (1S\*, 2S\*, 4S\*, 5R\*, 6S\*) 2-Amino-4-(4-chloro-benzoylamino)-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid




The title compound was prepared utilizing the two step procedure in Example 22. In step (a) the product of Preparation 1 (0.34 g, 0.95 mmol) and 4-chlorobenzoyl chloride (0.2 g, 1.15 mmol) were reacted to afford (1S\*, 2S\*, 4S\*, 5R\*, 6S\*)-2-tert-butoxycarbonylamino-4-(4-chloro-benzoylamino)-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid diethyl ester in 86.9% yield. In step (b) the intermediate amide (0.4 g, 0.81 mmol) was converted to the title compound and isolated as a white solid in a 87.5% yield (0.24 g, 0.71 mmol) by isoelectric precipitation ( pH = ~3-4 ).

m.p. > 250°C.

FDMS: M<sup>+</sup>+1 = 339.

Example 26

Synthesis of (1S\*, 2S\*, 4S\*, 5R\*, 6S\*) 2-Amino-4-(3-chloro-benzoylamino)-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid



The title compound was prepared utilizing the two step procedure in Example 22. In step (a) the product of Preparation 1 (0.33 g, 0.93 mmol) and 3-chlorobenzoyl chloride (0.19 g, 1.11 mmol) were reacted to afford (1S\*, 2S\*, 4S\*, 5R\*, 6S\*)-2-tert-butoxycarbonylamino-4-(3-chloro-benzoylamino)-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid diethyl ester in 99% yield. In step (b) the intermediate amide (0.45 g, 0.91 mmol) was converted to the title compound and isolated as a white solid in a 67.6% yield (0.208 g, 0.61 mmol) by isoelectric precipitation (pH = ~3-4).

m.p. = 254-255°C.


FDMS:  $M^{+}+1 = 339$ .

Anal. calcd. For  $C_{15}H_{15}N_2O_5Cl \cdot 0.1 H_2O$ : C, 52.90; H, 4.50; N, 8.23.

Found: C, 52.88; H, 4.40; N, 8.16.

Example 27

Synthesis of (1S\*, 2S\*, 4S\*, 5R\*, 6S\*) 2-Amino-4-(4-methoxybenzoylamino)-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid



The title compound was prepared utilizing the two step procedure in Example 22. In step (a) the product of Preparation 1 (0.3 g, 0.84 mmol) and 3-methoxybenzoyl chloride (0.17 g, 1.01 mmol) were reacted to afford (1S\*, 2S\*, 4S\*, 5R\*, 6S\*)-2-tert-butoxycarbonylamino-4-(4-methoxybenzoylamino)-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid diethyl ester in 77.7% yield. In step (b) the intermediate amide (0.3 g, 0.61 mmol) was converted to the title compound in a 64.6% yield (0.132 g, 0.4 mmol).

m.p. = 248-249°C.


FDMS:  $M^+ + 1 = 335$ .

Anal. calcd. For  $C_{16}H_{18}N_2O_6 \cdot 0.3 H_2O$ : C, 56.57; H, 5.52; N, 8.25.

Found: C, 56.67; H, 5.25; N, 8.22.

Example 28

Synthesis of (1S\*, 2S\*, 4S\*, 5R\*, 6S\*) 2-Amino-4-[(naphthalene-1-carbonyl)-amino]-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid




The title compound was prepared utilizing the two step procedure in Example 22. In step (a) the product of Preparation 1 (0.33 g, 0.93 mmol) and 1-naphthoyl chloride (0.212 g, 1.1 mmol) were reacted to afford (1S\*, 2S\*, 4S\*, 5R\*, 6S\*)-2-tert-butoxycarbonylamino-4-[(naphthalene-1-carbonyl)-amino]-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid diethyl ester in 99% yield. In step (b) the intermediate amide (0.39 g, 0.76 mmol) was converted to the title compound in a 51.7% yield (0.14 g, 0.4 mmol).

m.p. = 234-235°C.

FDMS: M<sup>+</sup>+1 = 355.

Example 29

Synthesis of (1S\*, 2S\*, 4S\*, 5R\*, 6S\*) 2-Amino-4-[(naphthalene-2-carbonyl)-amino]-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid



The title compound was prepared utilizing the two step procedure in Example 22. In step (a) the product of Preparation 1 (0.33 g, 0.93 mmol) and 2-naphthoyl chloride (0.21 g, 1.1 mmol) were reacted to afford (1S\*, 2S\*, 4S\*, 5R\*, 6S\*)-2-tert-butoxycarbonylamino-4-[(naphthalene-2-carbonyl)-amino]-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid diethyl ester in 99% yield. In step (b) the intermediate amide (0.43 g, 0.84 mmol) was converted to the title compound and isolated as a white solid in an 88.7% yield (0.264 g, 0.745 mmol) by isoelectric precipitation (pH = ~3-4).

m.p. = 257-258°C.

FDMS:  $M^{+}+1$  = 355.


Anal. calcd. For  $C_{19}H_{18}N_2O_5 \cdot 0.2 H_2O$ : C, 63.75; H, 5.80; N, 7.83.

Found: C, 63.79; H, 4.96; N, 7.78.



Example 30

Synthesis of (1S\*, 2S\*, 4S\*, 5R\*, 6S\*) 2-Amino-4-(4-fluorobenzoylamino)-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid



The title compound was prepared utilizing the two step procedure in Example 22. In step (a) the product of Preparation 1 (0.15 g, 0.42 mmol) and 4-fluorobenzoyl chloride (0.08 g, 0.5 mmol) were reacted to afford (1S\*, 2S\*, 4S\*, 5R\*, 6S\*)-2-tert-butoxycarbonylamino-4-(4-fluorobenzoylamino)-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid diethyl ester in 89.4% yield. In step (b) the intermediate amide (0.15 g, 0.31 mmol) was converted to the title compound and isolated as a solid in a 77.4% yield (0.077 g, 0.24mmol) by isoelectric precipitation (pH = ~3-4).

m.p. > 250°C.


FDMS:  $M^+ + 1 = 323$ .

Anal. calcd. For  $C_{15}H_{15}N_2O_5F \cdot 0.5 H_2O$ : C, 54.38; H, 4.87; N, 8.46.

Found: C, 54.46; H, 4.61; N, 8.41.

Example 31

Synthesis of (1S\*, 2S\*, 4S\*, 5R\*, 6S\*) 2-Amino-4-[(thiophene-2-carbonyl)-amino]-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid



The title compound was prepared utilizing the two step procedure in Example 22. In step (a) the product of Preparation 1 (0.15 g, 0.42 mmol) and 2-thiophenecarbonyl chloride (0.074 g, 0.51 mmol) were reacted to afford (1S\*, 2S\*, 4S\*, 5R\*, 6S\*)-2-tert-butoxycarbonylamino-4-[(thiophene-2-carbonyl)-amino]-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid diethyl ester in 79% yield. In step (b) the intermediate amide (0.14 g, 0.3 mmol) was converted to the title compound in a 49.4% yield (0.046 g, 0.16 mmol).

m.p. = 254-255°C.


FDMS:  $M^+$  = 310.

Anal. calcd. For  $C_{13}H_{14}N_2O_5S \cdot 2.0 H_2O$ : C, 45.08; H, 5.24; N, 8.09.

Found: C, 45.05; H, 5.00; N, 8.09.

Example 32

Synthesis of (1S\*, 2S\*, 4S\*, 5R\*, 6S\*) 2-Amino-4-(2-methoxybenzoylamino)-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid



The title compound was prepared utilizing the two step procedure in Example 22. In step (a) the product of Preparation 1 (0.16 g, 0.45 mmol) and 2-methoxybenzoyl chloride (0.092 g, 0.54 mmol) were reacted to afford (1S\*, 2S\*, 4S\*, 5R\*, 6S\*)-2-tert-butoxycarbonylamino-4-(2-methoxybenzoylamino)-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid diethyl ester in 86.1% yield. In step (b) the intermediate amide (0.15 g, 0.31 mmol) was converted to the title compound in a 40.9% yield (0.045 g, 0.14 mmol).

m.p. = 247-248°C.


FDMS:  $M^+ + 1 = 335$ .

Anal. calcd. For  $C_{16}H_{18}N_2O_6 \cdot 0.7 H_2O$ : C, 55.39; H, 5.64; N, 8.08.

Found: C, 55.17; H, 5.29; N, 8.00.

Example 33

Synthesis of (1S\*, 2S\*, 4S\*, 5R\*, 6S\*) 2-Amino-4-(3-methoxybenzoylamino)-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid



The title compound was prepared utilizing the two step procedure in Example 22. In step (a) the product of Preparation 1 (0.16 g, 0.45 mmol) and 3-methoxybenzoyl chloride (0.092 g, 0.54 mmol) were reacted to afford (1S\*, 2S\*, 4S\*, 5R\*, 6S\*)-2-tert-butoxycarbonylamino-4-(3-methoxybenzoylamino)-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid diethyl ester in 95.1% yield. In step (b) the intermediate amide (0.19 g, 0.39 mmol) was converted to the title compound in a 54.1% yield (0.070 g, 0.21 mmol).

m.p. = 235-236°C.


FDMS:  $M^{+}+1$  = 335.

Anal. calcd. For  $C_{16}H_{18}N_2O_6 \cdot 1.2 H_2O$ : C, 53.99; H, 5.78; N, 7.87.

Found: C, 54.01; H, 5.45; N, 7.87.

Example 34

Synthesis of (1S\*, 2S\*, 4S\*, 5R\*, 6S\*) 2-Amino-4-(3-methyl-benzoylamino)-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid



The title compound was prepared utilizing the two step procedure in Example 22. In step (a) the product of Preparation 1 (0.2 g, 0.56 mmol) and m-toluoyl chloride (0.104 g, 0.67 mmol) were reacted to afford (1S\*, 2S\*, 4S\*, 5R\*, 6S\*)-2-tert-butoxycarbonylamino-4-(3-methyl-benzoylamino)-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid diethyl ester in 86.5% yield. In step (b) the intermediate amide (0.2 g, 0.42 mmol) was converted to the title compound in a 44.9% yield (0.060 g, 0.19 mmol).

m.p. > 250°C.


FDMS:  $M^{+}+1 = 319$ .

Anal. calcd. For  $C_{16}H_{18}N_2O_5 \cdot 0.2 H_2O$ : C, 59.69; H, 5.76; N, 8.70.

Found: C, 59.55; H, 5.53; N, 8.53.

Example 35

Synthesis of (1S\*, 2S\*, 4S\*, 5R\*, 6S\*) 2-Amino-4-(3-trifluoromethyl-benzoylamino)-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid



The title compound was prepared utilizing the two step procedure in Example 1. In step (a) the product of Preparation 1 (0.25 g, 0.7 mmol) and 3-(trifluoromethyl)benzoyl chloride (0.175 g, 0.84 mmol) was reacted to afford (1S\*, 2S\*, 4S\*, 5R\*, 6S\*)-2-tert-butoxycarbonylamino-4-(3-trifluoromethyl-benzoylamino)-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid diethyl ester in 99% yield. In step (b) the intermediate amide (0.26 g, 0.49 mmol) was converted to the title compound and isolated as a solid in a 67.4% yield (0.123 g, 0.33 mmol) by isoelectric precipitation (pH = ~3-4).

m.p. = 249-250°C.


FDMS:  $M^+ + 1 = 373$ .

Anal. calcd. For  $C_{16}H_{15}N_2O_5F_3 \cdot 0.2 H_2O$ : C, 51.12; H, 4.13; N, 7.45.

Found: C, 50.94; H, 3.99; N, 7.41.

Example 36

Synthesis of (1S\*, 2S\*, 4S\*, 5R\*, 6S\*) 2-Amino-4-(3-fluorobenzoylamino)-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid



The title compound was prepared utilizing the two step procedure in Example 22. In step (a) the product of Preparation 1 (0.23 g, 0.65 mmol) and 3-fluorobenzoyl chloride (0.1 g, 0.77 mmol) were reacted to afford (1S\*, 2S\*, 4S\*, 5R\*, 6S\*)-2-tert-butoxycarbonylamino-4-(3-fluorobenzoylamino)-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid diethyl ester 80% yield. In step (b) the intermediate amide (0.22 g, 0.46 mmol) was converted to the title compound and isolated as a solid in a 58.0% yield (0.086 g, 0.27 mmol) by isoelectric precipitation (pH = ~3-4).

m.p. > 250°C.


FDMS:  $M^{+}-1 = 321$ .

Anal. calcd. For  $C_{15}H_{15}N_2O_5F \cdot 0.3 H_2O$ : C, 54.98; H, 4.80; N, 8.55.

Found: C, 54.92; H, 4.63; N, 8.48.

Example 37

Synthesis of (1S\*, 2S\*, 4S\*, 5R\*, 6S\*) 2-Amino-4-(3,4-dichloro-benzoylamino)-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid



The title compound was prepared utilizing the two step procedure in Example 22. In step (a) the product of Preparation 1 (0.26 g, 0.73 mmol) and 3,4-dichlorobenzoyl chloride (0.183 g, 0.88 mmol) were reacted to afford (1S\*, 2S\*, 4S\*, 5R\*, 6S\*)-2-tert-butoxycarbonylamino-4-(3,4-dichloro-benzoylamino)-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid diethyl ester in 95.7% yield. In step (b) the intermediate amide (0.28 g, 0.53 mmol) was converted to the title compound and isolated as a solid in a 73.8% yield (0.146 g, 0.39 mmol) by isoelectric precipitation (pH = ~3-4).

m.p. > 250°C.

FDMS:  $M^+$  = 373.


Anal. calcd. For  $C_{15}H_{14}N_2O_5Cl_2 \cdot 0.09 C_3H_8O$ : C, 48.44; H, 3.91; N, 7.39.

Found: C, 48.77; H, 3.51; N, 7.49.



Example 38

Synthesis of (1S\*, 2S\*, 4S\*, 5R\*, 6S\*) 2-Amino-4-(3,5-dichloro-benzoylamino)-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid



The title compound was prepared utilizing the two step procedure in Example 22. In step (a) the product of Preparation 1 (0.22 g, 0.62 mmol) and 3,5-dichlorobenzoyl chloride (0.155 g, 0.74 mmol) were reacted to afford (1S\*, 2S\*, 4S\*, 5R\*, 6S\*)-2-tert-Butoxycarbonylamino-4-(3,5-dichloro-benzoylamino)-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid diethyl ester in 91% yield. In step (b) the intermediate amide (0.285 g, 0.54 mmol) was converted to the title compound and isolated as a solid in a 84.6% yield (0.17 g, 0.46 mmol) by isoelectric precipitation (pH = ~3-4).

m.p. = 250-251°C.


FDMS:  $M^+$  = 373.

Anal. calcd. For  $C_{15}H_{14}N_2O_5Cl_2$ : C, 48.28; H, 3.78; N, 7.33.

Found: C, 48.02; H, 3.78; N, 7.33.

Example 39

Synthesis of (1S\*, 2S\*, 4S\*, 5R\*, 6S\*) 2-Amino-4-(2,4-dichloro-benzoylamino)-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid



The title compound was prepared utilizing the two step procedure in Example 22. In step (a) the product of Preparation 1 (0.24 g, 0.67 mmol) and 2,4-dichlorobenzoyl chloride (0.169 g, 0.81 mmol) were reacted to afford (1S\*, 2S\*, 4S\*, 5R\*, 6S\*)-2-tert-butoxycarbonylamino-4-(2,4-dichloro-benzoylamino)-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid diethyl ester in 87% yield. In step (b) the intermediate amide (0.31 g, 0.59 mmol) was converted to the title compound and isolated as a solid in a 67.3% yield (0.147 g, 0.39 mmol) by isoelectric precipitation (pH = ~3-4).

m.p. = 231-232°C.


FDMS: M<sup>+</sup> = 373.

Anal. calcd. For C<sub>15</sub>H<sub>14</sub>N<sub>2</sub>O<sub>5</sub>Cl<sub>2</sub>.: C, 45.63; H, 4.19; N, 7.10.

Found: C, 45.40; H, 3.96; N, 7.47.

Example 40

Synthesis of (1S\*, 2S\*, 4S\*, 5R\*, 6S\*) 2-Amino-4-(2,3-dichloro-benzoylamino)-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid



The title compound was prepared utilizing the two step procedure in Example 22. In step (a) the product of Preparation 1 (0.23 g, 0.65 mmol) and 2,3-dichlorobenzoyl chloride (0.162 g, 0.77 mmol) were reacted to afford (1S\*, 2S\*, 4S\*, 5R\*, 6S\*)-2-tert-Butoxycarbonylamino-4-(2,3-dichloro-benzoylamino)-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid diethyl ester in 81% yield. In step (b) the intermediate amide (0.275 g, 0.52 mmol) was converted to the title compound and isolated as a solid in a 70.6% yield (0.137 g, 0.37 mmol) by isoelectric precipitation (pH = ~3-4).

m.p. = 249-250°C.


FDMS:  $M^+$  = 373.

Anal. calcd. For  $C_{15}H_{14}N_2O_5Cl_2 \cdot 1.0 H_2O$ : C, 46.05; H, 4.12; N, 7.16.

Found: C, 46.19; H, 3.92; N, 7.12.

Example 41

Synthesis of (1S\*, 2S\*, 4S\*, 5R\*, 6S\*) 2-Amino-4-(2-fluorobenzoylamino)-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid



The title compound was prepared utilizing the two step procedure in Example 22. In step (a) the product of Preparation 1 (0.245 g, 0.69 mmol) and 2-fluorobenzoyl chloride (0.131 g, 0.83 mmol) were reacted to afford (1S\*, 2S\*, 4S\*, 5R\*, 6S\*)-2-tert-butoxycarbonylamino-4-(2-fluorobenzoylamino)-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid diethyl ester in 78% yield. In step (b) the intermediate amide (0.25 g, 0.52 mmol) was converted to the title compound and isolated as a solid in a 62.1% yield (0.104 g, 0.32 mmol) by isoelectric precipitation (pH = ~3-4).

m.p. = □C.


FDMS:  $M^+ - 1 = 321$ .

Anal. calcd. For  $C_{15}H_{15}N_2O_5F \cdot 1.4 H_2O$ : C, 51.84; H, 5.16; N, 8.06.

Found: C, 51.61; H, 4.81; N, 8.11.

Example 42

Synthesis of (1S\*, 2S\*, 4S\*, 5R\*, 6S\*) 2-Amino-4-[(1H-indole-2-carbonyl)-amino]-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid



The title compound was prepared utilizing the two step procedure in Example 23. In step (a) the product of Preparation 1 (0.25 g, 0.7 mmol) and 2-indolecarboxylic acid (0.123 g, 0.77 mmol) were reacted to afford (1S\*, 2S\*, 4S\*, 5R\*, 6S\*)-2-tert-butoxycarbonylamino-4-[(1H-indole-2-carbonyl)-amino]-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid diethyl ester in 91.5% yield. In step (b) the intermediate amide (0.3 g, 0.6 mmol) was converted to the title compound in a 68.9% yield (0.137 g, 0.39 mmol).

m.p. > 250°C.


FDMS:  $M^+ - 1 = 342$ .

Anal. calcd. For  $C_{17}H_{17}N_3O_5 \cdot 0.3 H_2O$ : C, 58.55; H, 5.09; N, 12.05.

Found: C, 58.65; H, 4.86; N, 11.88.

Example 43

Synthesis of (1S\*, 2S\*, 4S\*, 5R\*, 6S\*) 2-Amino-4-(4-hydroxy-benzoylamino)-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid



The title compound was prepared utilizing the two step procedure in Example 23. In step (a) the product of Preparation 1 (0.25 g, 0.7 mmol) and 4-hydroxybenzoic acid (0.106 g, 0.77 mmol) were reacted to afford (1S\*, 2S\*, 4S\*, 5R\*, 6S\*)-2-tert-butoxycarbonylamino-4-(4-hydroxy-benzoylamino)-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid diethyl ester in 93% yield. In step (b) the intermediate amide (0.27 g, 0.56 mmol) was converted to the title compound in a 72.5% yield (0.131 g, 0.41 mmol).

m.p. = 252-253°C.


FDMS:  $M^+ - 1 = 319$ .

Anal. calcd. For  $C_{15}H_{16}N_2O_6 \cdot 1.1 H_2O$ : C, 52.97; H, 5.39; N, 8.24.

Found: C, 52.84; H, 5.02; N, 8.20.

Example 44

Synthesis of (1S\*, 2S\*, 4S\*, 5R\*, 6S\*) 2-Amino-4-(3-hydroxy-benzoylamino)-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid



The title compound was prepared utilizing the two step procedure in Example 23. In step (a) the product of Preparation 1 (0.25 g, 0.7 mmol) and 3-hydroxybenzoic acid (0.106 g, 0.77 mmol) were reacted to afford (1S\*, 2S\*, 4S\*, 5R\*, 6S\*)-2-tert-butoxycarbonylamino-4-(3-hydroxy-benzoylamino)-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid diethyl ester in 93% yield. In step (b) the intermediate amide (0.31 g, 0.65 mmol) was converted to the title compound in a 62.9% yield (0.131 g, 0.41 mmol).

m.p. > 250°C.


FDMS:  $M^+ + 1 = 321$ .

Anal. calcd. For  $C_{15}H_{16}N_2O_6 \cdot 0.4 H_2O$ : C, 55.01; H, 5.17; N, 8.55.

Found: C, 54.97; H, 5.00; N, 8.52.

Example 45

Synthesis of (1S\*, 2S\*, 4S\*, 5R\*, 6S\*) 2-Amino-4-(2-hydroxy-benzoylamino)-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid



The title compound was prepared utilizing the two step procedure in Example 23. In step (a) the product of Preparation 1 (0.25 g, 0.7 mmol) and 2-hydroxybenzoic acid (0.106 g, 0.77 mmol) were reacted to afford (1S\*, 2S\*, 4S\*, 5R\*, 6S\*)-2-tert-butoxycarbonylamino-4-(2-hydroxy-benzoylamino)-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid diethyl ester in 76% yield. In step (b) the intermediate amide (0.22 g, 0.46 mmol) was converted to the title compound in a 78.5% yield (0.116 g, 0.36 mmol).

m.p. > 250°C.

FDMS:  $M^{+}+1 = 321$ .


Anal. calcd. For  $C_{15}H_{16}N_2O_6 \cdot 0.55 H_2O$ : C, 54.71; H, 5.20; N, 8.51.

Found: C, 54.69; H, 4.88; N, 8.82.



Example 46

Synthesis of (1S\*, 2S\*, 4S\*, 5R\*, 6S\*) 2-Amino-4-[(biphenyl-4-carbonyl)-amino]-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid



The title compound was prepared utilizing the two step procedure in Example 23. In step (a) the product of Preparation 1 (0.25 g, 0.7 mmol) and 4-biphenylcarboxylic acid (0.153 g, 0.77 mmol) were reacted to afford (1S\*, 2S\*, 4S\*, 5R\*, 6S\*)-4-[(biphenyl-4-carbonyl)-amino]-2-tert-butoxycarbonylamino-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid diethyl ester in 98.5% yield. In step (b) the intermediate amide (0.36 g, 0.67 mmol) was converted to the title compound in a 76.5% yield (0.195 g, 0.51 mmol).

m.p. = 260-261°C.


FDMS:  $M^+ + 1 = 381$ .

Anal. calcd. For  $C_{22}H_{20}N_2O_6$ : C, 66.31; H, 5.30; N, 7.36.

Found: C, 66.00; H, 5.40; N, 7.38.

Example 47

Synthesis of (1S\*, 2S\*, 4S\*, 5R\*, 6S\*) 2-Amino-4-[(1H-indole-3-carbonyl)-amino]-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid



The title compound was prepared utilizing the two step procedure in Example 23. In step (a) the product of Preparation 1 (0.25 g, 0.7 mmol) and indole-3-carboxylic acid (0.124 g, 0.77 mmol) were reacted to afford (1S\*, 2S\*, 4S\*, 5R\*, 6S\*)-2-tert-butoxycarbonylamino-4-[(1H-indole-3-carbonyl)-amino]-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid diethyl ester in 48.6% yield. In step (b) the intermediate amide (0.16 g, 0.32 mmol) was converted to the title compound in a 91.9% yield (0.101 g, 0.29 mmol).

m.p. = 245-246°C.


FDMS:  $M^+ + 1 = 344$ .

Anal. calcd. For  $C_{17}H_{17}N_3O_5$ : C, 56.47; H, 4.88; N, 11.62.

Found: C, 56.52; H, 4.85; N, 11.48.

Example 48

Synthesis of (1S\*, 2S\*, 4S\*, 5R\*, 6S\*) 2-Amino-4-[(isoquinoline-1-carbonyl)-amino]-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid



The title compound was prepared utilizing the two step procedure in Example 23. In step (a) the product of Preparation 1 (0.25 g, 0.7 mmol) and 1-isoquinolinecarboxylic acid (0.133 g, 0.77 mmol) were reacted to afford (1S\*, 2S\*, 4S\*, 5R\*, 6S\*)-2-tert-butoxycarbonylamino-4-[(isoquinoline-1-carbonyl)-amino]-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid diethyl ester 89.4% yield. In step (b) the intermediate amide (0.3 g, 0.59 mmol) was converted to the title compound and isolated as a solid in 56.8% yield (0.119 g, 0.34 mmol) by cation exchange chromatography (Dowex 50X8-100: 10% Pyridine/H<sub>2</sub>O).

m.p. = 225-226°C.


FDMS: M<sup>+</sup>+1= 356.

Anal. calcd. For C<sub>18</sub>H<sub>17</sub>N<sub>3</sub>O<sub>5</sub>·0.7 H<sub>2</sub>O: C, 58.75; H, 5.04; N, 11.42.

Found: C, 58.43; H, 4.73; N, 11.67.

Example 49

Synthesis of (1S\*, 2S\*, 4S\*, 5R\*, 6S\*)-2-Amino-4-[(pyridine-2-carbonyl)-amino]-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid



The title compound was prepared by the two step procedure in Example 1. The product of Preparation 1 (0.30g, 0.84 mmol) was reacted with picolinoyl chloride hydrochloride (0.18g, 1.02 mmol) in step (a). Purification of the desired intermediate was by PC-TLC (13% EtOAc/Hexane to 100% EtOAc) with a 90% yield. The intermediate, (1S\*, 2S\*, 4S\*, 5R\*, 6S\*)-2-tert-Butoxycarbonylamino-4-[(pyridine-2-carbonyl)-amino]-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid diethyl ester (0.34g, 0.74 mmol) was converted to the title compound by step(b) using isoelectric point crystalization from H<sub>2</sub>O, pH 3, in a 84% yield (0.19g, 0.62 mmol).

m.p.=228-229°C.


FDMS: M<sup>+</sup>+1=306.

Anal. Calcd. For C<sub>14</sub>H<sub>15</sub>N<sub>3</sub>O<sub>5</sub>.0.4 H<sub>2</sub>O: C, 53.81; H, 5.10; N, 13.45.

Found: C, 53.94; H, 5.10; N, 13.31.

Example 50

Synthesis of (1S\*, 2S\*, 4S\*, 5R\*, 6S\*)-2-Amino-4-[(pyridine-3-carbonyl)-amino]-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid



The title compound was prepared by the two step procedure in Example 1. The product of Preparation 1 (0.30g, 0.84 mmol) was reacted with nicotinoyl chloride hydrochloride (0.18g, 1.01 mmol) in step (a). Purification of the desired intermediate was by PC-TLC (10% EtOAc/Hexane to 100% EtOAc) with a 90% yield. The intermediate, (1S\*, 2S\*, 4S\*, 5R\*, 6S\*)-2-*tert*-Butoxycarbonylamino-4-[pyridine-3-carbonyl]-amino]-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid diethyl ester (0.35g, 0.76 mmol) was converted into the title compound by step (b) using isoelectric point crystallization from H<sub>2</sub>O, pH 3, in a 53% yield (0.12g, 0.40 mmol).

m.p.=223-224°C.


FDMS: M<sup>+</sup>+1=306.

Anal. Calcd. For C<sub>14</sub>H<sub>15</sub>N<sub>3</sub>O<sub>5</sub>.1.0 H<sub>2</sub>O: C, 52.01; H, 5.30; N, 13.00.

Found: C, 51.77; H, 5.18; N, 12.74.

Example 51

Synthesis of (1S\*, 2S\*, 4S\*, 5R\*, 6S\*)-2-Amino-4-[(pyridine-4-carbonyl)-amino]-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid



The title compound was prepared by the two step procedure in Example 1. The product of Preparation 1 (0.30g, 0.84 mmol) was reacted with isonicotinoyl chloride hydrochloride (0.18g, 1.01 mmol) in step (a). Purification of the desired intermediate was by PC-TLC (25% EtOAc/Hexane to 100% EtOAc) with a 80% yield. The intermediate, (1S\*, 2S\*, 4S\*, 5R\*, 6S\*)-2-tert-Butoxycarbonylamino-4-[pyridine-4-carbonyl]-amino]-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid diethyl ester (0.31g, 0.68 mmol) was converted into the title compound by step (b) using isoelectric point crystallization from H<sub>2</sub>O, pH 3, in a 37% yield (0.76g, 0.25 mmol).

m.p. => 260°C.


FDMS:  $M^+ - 1 = 304$ .

Anal. Calcd. For  $C_{14}H_{15}N_3O_5 \cdot 0.6 HCl$ : C, 55.08; H, 4.95; N, 13.76.

Found: C, 51.27; H, 4.80; N, 12.59.

Example 52

Synthesis of (1S\*, 2S\*, 4S\*, 5R\*, 6S\*)-2-Amino-4-(2-trifluoromethyl-benzoylamino)-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid



The title compound was prepared by the two step procedure in Example 1. The product of Preparation 1 (0.30g, 0.84 mmol) was reacted with 2-(trifluoro-methyl)benzoyl chloride (0.21g, 1.01 mmol) in step (a). Purification of the desired intermediate was by PC-TLC (10% EtOAc/Hexane to 50% EtOAc/Hexane) with a 99% yield (0.44g, 0.83 mmol). The intermediate, (1S\*, 2S\*, 4S\*, 5R\*, 6S\*)-2-*tert*-Butoxycarbonylamino-4-(2-trifluoromethyl-benzoylamino)-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid diethyl ester (0.42g, 0.80 mmol) was converted into the title compound by step(b) using isoelectric point crystallization from H<sub>2</sub>O, pH 3, in a 63% yield (0.19g, 0.50 mmol).

m.p.=243-244°C.

FDMS: M<sup>+</sup>+1=373.


Anal. Calcd. For C<sub>16</sub>H<sub>15</sub>N<sub>2</sub>O<sub>5</sub>F<sub>3</sub>.0.5 H<sub>2</sub>O: C, 50.40; H, 4.23; N, 7.35.

Found: C, 50.04; H, 3.83; N, 7.29.

-111-

Example 53

Synthesis of (1S\*, 2S\*, 4S\*, 5R\*, 6S\*)-2-Amino-4-(3,5-difluoro-benzoylamino)-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid



The title compound was prepared by the two step procedure in Example 1. The product of Preparation 1 (0.30g, 0.84 mmol) was reacted with 3,5-difluorobenzoyl chloride (0.18g, 1.01 mmol) in step (a). Purification of the desired intermediate was by PC-TLC (10% EtOAc/Hexane to 50% EtOAc/Hexane) with a 67% yield. The intermediate, (1S\*, 2S\*, 4S\*, 5R\*, 6S\*)-2-tert-Butoxycarbonylamino-4-(3,5-difluoro-benzoylamino)-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid diethyl ester (0.28g, 0.56 mmol) was converted into the title compound by step (b) using isoelectric point crystalization from H<sub>2</sub>O, pH 2-3 in a 83% yield (0.16g, 0.47 mmol).

m.p. >260°C.

FDMS: M<sup>+</sup>+1=341.

Anal. Calcd. For C<sub>15</sub>H<sub>14</sub>N<sub>2</sub>O<sub>5</sub>F<sub>2</sub>.0.1 H<sub>2</sub>O: C, 52.67; H, 4.18; N, 8.19.


Found: C, 52.28; H, 3.93; N, 8.03.



-112-

Example 54

Synthesis of (1S\*, 2S\*, 4S\*, 5R\*, 6S\*)-2-Amino-4-(2-methyl-benzoylamino)bicyclo[3.1.0]hexane-2,6-dicarboxylic acid



The title compound was prepared by the two step procedure in Example 1. The product of Preparation 1 (0.30g, 0.84 mmol) was reacted with o-toluoyl chloride. (0.16g, 1.01 mmol) in step (a). Purification of the desired intermediate was by PC-TLC (10% EtOAc/Hexane to 50% EtOAc/Hexane) with a 90% yield (0.36g, 0.76 mmol). The intermediate, (1S\*, 2S\*, 4S\*, 5R\*, 6S\*)-2-*tert*-Butoxycarbonylamino-4-(2-methyl-benzoylamino)-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid diethyl ester (0.34g, 0.72 mmol) was converted into the title compound by step (b) using isoelectric point crystalization from H<sub>2</sub>O, pH 3, in a 64% yield (0.15g, 0.46 mmol).

m.p.=228-229°C.


FDMS: M<sup>+</sup>+1=319.

Anal. Calcd. For C<sub>16</sub>H<sub>18</sub>N<sub>2</sub>O<sub>5</sub>·0.5 H<sub>2</sub>O: C, 58.71; H, 5.85; N, 8.56.

Found: C, 58.49; H, 6.47; N, 8.47.

Example 55

Synthesis of (1S\*, 2S\*, 4S\*, 5R\*, 6S\*)-2-Amino-4-(4-methyl-benzoylamino)-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid



The title compound was prepared by the two step procedure in Example 1. The product of Preparation 1 (0.30g, 0.84 mmol) was reacted with *p*-toluoyl chloride (0.16g, 1.01 mmol) in step(a). Purification of the desired intermediate was by PC-TLC (10% EtOAc/Hexane to 30% EtOAc/Hexane) with a 95% yield (0.38g, 0.80 mmol). The intermediate, (1S\*, 2S\*, 4S\*, 5R\*, 6S\*)-2-*tert*-Butoxycarbonylamino-4-(4-methyl-benzoylamino)-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid diethyl ester (0.37g, 0.77 mmol) was converted into the title compound by step (b) using isoelectric point crystalization from H<sub>2</sub>O, pH 3, in a 89% yield (0.22g, 0.69 mmol).

m.p.=243-244°C.


FDMS: M<sup>+</sup>+1=319.

Anal. Calcd. For C<sub>16</sub>H<sub>18</sub>N<sub>2</sub>O<sub>5</sub>·0.2 H<sub>2</sub>O: C, 59.69; H, 5.76; N, 8.70.

Found: C, 59.32; H, 5.57; N, 8.50.

Example 56

Synthesis of (1S\*, 2S\*, 4S\*, 5R\*, 6S\*)-2-Amino-4-[(quinoxaline-2-carbonyl)-amino]-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid



The title compound was prepared by the two step procedure in Example 1. The product of Preparation 1 (0.25g, 0.70 mmol) was reacted with 2-quinoxaloyl chloride (0.16g, 0.84 mmol) in step (a). Purification of the desired intermediate was by PC-TLC (25% EtOAc/Hexane to 100% EtOAc/Hexane) with a 84% yield (0.30g, 0.59 mmol). (1S\*, 2S\*, 4S\*, 5R\*, 6S\*)-2-*tert*-Butoxycarbonylamino-4-[(quinoxaline-2-carbonyl)-amino]-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid diethyl ester (0.29g, 0.57 mmol) was converted into the title compound by step (b) using isoelectric point crystalization from H<sub>2</sub>O, pH 3, in a 83% yield (0.17g, 0.47 mmol).

m.p. >260°C.


FDMS: M<sup>+</sup>+1=357.

Anal. Calcd. For C<sub>17</sub>H<sub>16</sub>N<sub>4</sub>O<sub>5</sub>·0.4 HCl: C, 55.05; H, 4.46; N, 15.11.

Found: C, 54.67; H, 4.22; N, 14.88.

Example 57

Synthesis of (1S\*, 2S\*, 4S\*, 5R\*, 6S\*)-2-Amino-4-(3-carboxy-benzoylamino)-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid



The title compound was prepared by the two step procedure in Example 2. The product of Preparation 1 (0.30g, 0.84 mmol) was reacted with monomethyl isophthalate (0.17g, 0.93 mmol) in step (a). Purification of the desired intermediate was by PC-TLC (10% EtOAc/Hexane to 30% EtOAc/Hexane) with a 78% yield (0.34g, 0.66 mmol). The intermediate, (1S\*, 2S\*, 4S\*, 5R\*, 6S\*)-2-*tert*-Butoxycarbonylamino-4-(3-carboxy-benzoylamino)-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid diethyl ester (0.31g, 0.60 mmol) was converted into the title compound by step (b) using isoelectric point crystalization from H<sub>2</sub>O, pH 3, in a 89% yield (0.19g, 0.53 mmol).

m.p.=249-250°C.


FDMS: M<sup>+</sup>+1=349.

Anal. Calcd. For C<sub>16</sub>H<sub>16</sub>N<sub>2</sub>O<sub>7</sub>·1.0 H<sub>2</sub>O: C, 52.46; H, 4.95; N, 7.65.

Found: C, 52.09; H, 4.59; N, 7.58.

Example 58

Synthesis of (1S\*, 2S\*, 4S\*, 5R\*, 6S\*)-2-Amino-4-(2-carboxy-benzoylamino)-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid



The title compound was prepared by the two step procedure in Example 2. The product of Preparation 1 (0.25g, 0.70 mmol) was reacted with monomethyl phthalate (0.14g, 0.77 mmol) in step (a). 1-[3-(Dimethylamino)propyl]-3-ethylcarbodiimide hydrochloride was used (0.20g, 1.05 mmol) in the place of 1,3-Dicyclohexylcarbodiimide. In the workup, extraction was EtOAc/H<sub>2</sub>O (pH=1), followed by a straight H<sub>2</sub>O wash of the organics. Purification of the desired intermediate was by PC-TLC (10% EtOAc/Hexane to 50% EtOAc/Hexane) with a 89% yield (0.32g, 0.62 mmol). The intermediate, (1S\*, 2S\*, 4S\*, 5R\*, 6S\*)-2-*tert*-Butoxycarbonylamino-4-(2-carboxy-benzoylamino)-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid diethyl ester (0.31g, 0.59 mmol) was converted into the title compound by step (b) using isoelectric point crystalization from H<sub>2</sub>O, pH 1-2, in a 68% yield (0.14g, 0.40 mmol).

m.p. >260°C.


FDMS: M<sup>+</sup>+1=349.

Anal. Calcd. For C<sub>16</sub>H<sub>16</sub>N<sub>2</sub>O<sub>7</sub>·0.5 HCl, 1.8 H<sub>2</sub>O: C, 48.17; H, 5.08; N, 7.02.

Found: C, 48.17; H, 4.80; N, 7.00.

Example 59

Synthesis of (1S\*, 2S\*, 4S\*, 5R\*, 6S\*)-2-Amino-4-(4-carboxy-benzoylamino)-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid



The title compound was prepared by the two step procedure in Example 2. The product of Preparation 1 (0.25g, 0.70 mmol) was reacted with monomethyl terephthalate (0.14g, 0.77 mmol) in step (a). 1-[3-(Dimethylamino)propyl]-3-ethylcarbodiimide hydrochloride was used (0.20g, 1.05 mmol) in the place of 1,3-Dicyclohexylcarbodiimide. In the workup, extraction was EtOAc/H<sub>2</sub>O (pH=1), followed by a straight H<sub>2</sub>O wash of the organics. Purification of the desired intermediate was by PC-TLC (10% EtOAc/Hexane to 30% EtOAc/Hexane) with a 88% yield (0.32g, 0.62 mmol). The intermediate, (1S\*, 2S\*, 4S\*, 5R\*, 6S\*)-2-tert-Butoxycarbonylamino-4-(4-carboxy-benzoylamino)-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid diethyl ester (0.31g, 0.59 mmol) was converted into the title compound by step (b) using isoelectric point crystallization from H<sub>2</sub>O, pH 1-2, in a 85% yield (0.18g, 0.50 mmol).

m.p. >260°C.


FDMS: M<sup>+</sup>+1=349.

Anal. Calcd. For C<sub>16</sub>H<sub>16</sub>N<sub>2</sub>O<sub>7</sub>·0.1 HCl, 0.7 H<sub>2</sub>O: C, 52.71; H, 4.84; N, 7.68.

Found: C, 52.52; H, 4.46; N, 7.52.

Example 60

Synthesis of (1S\*, 2S\*, 4S\*, 5R\*, 6S\*)-2-Amino-4-[(1 H-imidazole-4-carbonyl)-amino]-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid



The title compound was prepared by the two step procedure in Example 2. The product of Preparation 1 (0.25g, 0.70 mmol) was reacted with imidazol-4-carboxylic acid (0.0.9g, 0.77 mmol) in step (a). 1-[3-(Dimethylamino)propyl]-3-ethylcarboiimide hydrochloride was used (0.20g, 1.05 mmol) in the place of 1,3-Dicyclohexylcarbodiimide. In the workup, the organics were dried over anhydrous  $K_2CO_3$ . Purification of the desired intermediate was by PC-TLC (25% EtOAc/Hexane to 3% EtOH/EtOAc) with a 63% yield (0.20g, 0.44 mmol). The intermediate, (1S\*, 2S\*, 4S\*, 5R\*, 6S\*)-2-tert-Butoxycarbonylamino-4-[(1 H-imidazole-4-carbonyl)-amino]-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid diethyl ester (0.17g, 0.38 mmol) was converted into the title compound by step (b) using isoelectric point crystalization from  $H_2O$ , pH 3, in a 47% yield (0.05g, 0.18 mmol).

m.p.=258-259°C.


FDMS:  $M^+ + 1 = 295$ .

Anal. Calcd. For  $C_{12}H_{14}N_4O_5 \cdot 0.8 HCl$ : C, 44.56; H, 4.61; N, 17.32.

Found: C, 44.54; H, 4.65; N, 16.96.

Example 61

Synthesis of (1S\*, 2S\*, 4S\*, 5R\*, 6S\*)-2-Amino-4-(3-phenyl-acroloylamino)-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid



The title compound was prepared by the two step procedure in Example 2. The product of Preparation 1 (0.30g, 0.84 mmol) was reacted with trans-cinnamic acid (0.14g, 0.93 mmol) in step (a). 1-[3-(Dimethylamino)propyl]-3-ethylcarbodiimide hydrochloride was used (0.24g, 1.26 mmol) in the place of 1,3-Dicyclohexylcarbodiimide. In the workup, extraction was EtOAc/H<sub>2</sub>O (pH=1), followed by a straight H<sub>2</sub>O wash of the organics. Purification of the desired intermediate was by PC-TLC (10% EtOAc/Hexane to 30% EtOAc/Hexane) with a 63% yield (0.26g, 0.53 mmol). The intermediate, (1S\*, 2S\*, 4S\*, 5R\*, 6S\*)-2-tert-Butoxycarbonylamino-4-(3-phenyl-acroloylamino)-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid diethyl ester (0.26g, 0.53 mmol) was converted into the amine by step (b), the Boc group removal, 99.9% yield (0.21g, 0.54 mmol). FDMS: M<sup>+</sup>+1=387. After hydrolysis of the esters, the title compound was isolated using isoelectric point crystalization from H<sub>2</sub>O, pH 3, in a 79% yield (0.13g, 0.39 mmol).

m.p.=225-226°C.

FDMS: M<sup>+</sup>+1=331.


Anal. Calcd. For C<sub>17</sub>H<sub>18</sub>N<sub>2</sub>O<sub>5</sub>. C, 61.81; H, 5.49; N, 8.48.

Found: C, 61.52; H, 5.65; N, 8.12.



Example 62

Synthesis of (1S\*, 2S\*, 4S\*, 5R\*, 6S\*)-2-Amino-4-(3-phenyl-propynoylamino)-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid



The title compound was prepared by the two step procedure in Example 2. The product of Preparation 1 (0.25g, 0.70 mmol) was reacted with phenylpropionic acid (0.11g, 0.77 mmol) in step (a). 1-[3-(Dimethylamino)propyl]-3-ethylcarbodiimide hydrochloride was used (0.20g, 1.05 mmol) in the place of 1,3-Dicyclohexylcarbodiimide. In the workup, extraction was EtOAc/H<sub>2</sub>O (pH=1), followed by a straight H<sub>2</sub>O wash of the organics. Purification of the desired intermediate was by PC-TLC (10% EtOAc/Hexane to 30% EtOAc/Hexane) with a 48% yield (0.16g, 0.34 mmol). The intermediate, (1S\*, 2S\*, 4S\*, 5R\*, 6S\*)-2-*tert*-Butoxycarbonylamino-4-(3-phenyl-propynoylamino)-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid diethyl ester (0.15g, 0.31 mmol) was converted into the title compound by step (b) using isoelectric point crystalization from H<sub>2</sub>O, pH 3, in a 64% yield (0.07g, 0.20 mmol).

m.p.=222-223°C.


FDMS: M<sup>+</sup>+1=329.

Anal. Calcd. For C<sub>17</sub>H<sub>16</sub>N<sub>2</sub>O<sub>5</sub>·0.6 H<sub>2</sub>O: C, 60.21; H, 5.11; N, 8.26.

Found: C, 59.93; H, 5.05; N, 8.31.

Example 63

Synthesis of (1S\*, 2S\*, 4S\*, 5R\*, 6S\*)-2-Amino-4-benzylamino-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid



A. (1S\*, 2S\*, 4S\*, 5R\*, 6S\*)-4-Benzylamino-2-tert-butoxycarbonylamino-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid diethyl ester.

Product from Preparation 1 (a) (0.30g, 0.84 mmol) was dissolved in EtOH at ambient temperature under nitrogen. Benzyl bromide (0.15g, 0.91 mmol) was added in one portion. The reaction mixture was stirred for 20 hours. The reaction mixture was concentrated under vacuum. Purification by PC-TLC (14% EtOAc/hexanes to 100% EtOAc) afforded the title product as the desired intermediate (0.14g, 0.31 mmol).

FDMS: M<sup>+</sup> +1=447.

B. (1S\*, 2S\*, 4S\*, 5R\*, 6S\*)-2-Amino-4-benzylamino-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid.

(1S\*, 2S\*, 4S\*, 5R\*, 6S\*)-4-Benzylamino-2-tert-butoxycarbonylamino-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid diethylester from step (a) plus additional intermediate (0.24g, 0.54 mmol) in 30 mL of EtOAc was cooled to 0°C under nitrogen. HCl(g) was added until the solution was saturated. The reaction was concentrated under vacuum. THF

-122-

(8mL) and deionized H<sub>2</sub>O (1mL) were added to the reaction concentrate. 1N NaOH (9.5mL, 9.5 mmol) was added at ambient temperature, and the reaction was left stirring for 20 hour, pH 13-14. Using 1 N HCl, the reaction was taken to pH=7, concentrated under vacuum. The solids were reconstituted in H<sub>2</sub>O. Purification was by anion-exchange chromatography (Bio-Rad® AG1-X8: elution with 3N AcOH). Compound diluted in CH<sub>3</sub>OH, precipitate obtained upon addition of IPA. Crystals vacuum filtered, 80% yield (0.13g, 0.43 mmol).

mp=244-245°C.


FDMS: M<sup>+</sup>-1=289.

Anal. Calcd. For C<sub>15</sub>H<sub>18</sub>N<sub>2</sub>O<sub>4</sub>·0.1 IPA, 0.7 H<sub>2</sub>O: C, 59.48; H, 6.59; N, 9.07.

Found: C, 59.15; H, 6.39; N, 8.71.

Example 64

Synthesis of (1S\*,2S\*,4S\*,5R\*,6R\*) 2-Amino-4-phenylcarbamoyloxy-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid



A. (1S\*,2S\*,4S\*,5R\*,6R\*) Diethyl 2-(N-t-butyloxycarbonylamino)-4-phenylcarbamoyloxy-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid.

A 0°C solution of (1S\*,2S\*,4S\*,5R\*,6R\*) diethyl 2-(N-t-butyloxycarbonylamino)-4-hydroxybicyclo[3.1.0]hexane-2,6-dicarboxylic acid (0.30 g, 0.84 mmol—synthesis previously described in US 5,958,960 example 8a) in pyridine (10 mL) was treated in one portion with phenylisocyanate (0.12 g, 1.0 mmol) and allowed to warm to room temperature as it stirred overnight. The reaction mixture was partitioned between EtOAc and aqueous NaHSO<sub>4</sub> and the product extracted with EtOAc. All organic layers were combined, washed with brine, dried over MgSO<sub>4</sub>, concentrated in vacuo and purified by SiO<sub>2</sub> chromatography (PC-TLC: 10 % EtOAc/hexanes to 20 % EtOAc/hexanes) to yield 0.30 g (0.63 mmol, 75 %) of a white solid.

FDMS: M<sup>+</sup> + 1 = 477.

-124-

Anal. calcd. for  $C_{24}H_{32}N_2O_8 \cdot 0.1 \text{ eq. } CH_2Cl_2$ : C, 59.68; H, 6.69; N, 5.78.

Found: C, 59.75; H, 6.59; N, 5.68.

B. (1S\*,2S\*,4S\*,5R\*,6R\*) 2-Amino-4-phenylcarbamoyloxy-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid.

A 0 °C solution of the product of example 21a (0.28 g, 0.59 mmol) in EtOAc (35mL) was purged with anhydrous HCl gas until the solution was saturated. The resulting reaction mixture was stirred at 0 °C for 4 hours and concentrated to dryness in vacuo to afford the HCl salt. The resulting solids were reconstituted in a 1:1 mixture of 1N NaOH:THF (15 mL total volume) and stirred at room temperature overnight. The reaction mixture was adjusted to pH = 7 with 1N HCl and concentrated to dryness. The solids were reconstituted in water, adjusted to pH = 12 with 1N NaOH and purified by anion exchange chromatography (Bio-Rad® AG1-X8 resin; acetate form converted to hydroxide form). The product was eluted from the column with 3N AcOH and concentrated to dryness in vacuo to afford crude product. Trituration in 2-propanol/H<sub>2</sub>O (5:1) followed by drying under vacuum at 80 °C afforded 0.12 g (0.37 mmol, 63 %) of the desired product as a white solid.

mp = dec >270 °C.


FDMS:  $M^+ + 1 = 321$ .

Anal. calcd. for  $C_{15}H_{16}N_2O_6 \cdot 0.2 \text{ eq. } H_2O$ : C, 55.62; H, 5.10; N, 8.65.

Found: C, 55.51; H, 5.01; N, 8.39.

Example 65

Synthesis of (1S\*, 2S\*, 4S\*, 5R\*, 6R\*)-2-Amino-4-(2-chloro-phenylcarbamoyloxy)-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid



The title compound was prepared by the two step procedure Example Matt. (1S\*, 2S\*, 4S\*, 5R\*, 6R\*)-2-*tert*-Butoxycarbonylamino-4-hydroxy-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid diethyl ester (0.30g, 0.84 mmol) was reacted with *o*-chlorophenyl isocyanate (0.16g, 1.01 mmol) in step (a). Purification of the desired intermediate was by PC-TLC (10% EtOAc/Hexane to 25% EtOAc/Hexane) with a 59% yield (0.25g, 0.49 mmol). (1S\*, 2S\*, 4S\*, 5R\*, 6R\*)-2-*tert*-Butoxycarbonylamino-4-(2-chloro-phenylcarbamoyloxy)-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid diethyl ester (0.25g, 0.49 mmol) was converted into the title compound by step (b) using isoelectric point crystalization from H<sub>2</sub>O, pH 3, in a 37% yield (0.06g, 0.18 mmol).

m.p.=245-246°C.


FDMS: M<sup>+</sup>+1=355.

Anal. Calcd. For C<sub>15</sub>H<sub>15</sub>N<sub>2</sub>O<sub>6</sub>Cl.0.1 H<sub>2</sub>O: C, 50.53; H, 4.30; N, 7.86.

Found: C, 50.13; H, 3.93; N, 7.58.

Example 66

Synthesis of (1S\*, 2S\*, 4S\*, 5R\*, 6R\*)-2-Amino-4-(2-fluoro-phenylcarbamoyloxy)-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid



A. (1S\*, 2S\*, 4S\*, 5R\*, 6R\*)-2-tert-Butoxycarbonylamino-4-(2-fluoro-phenylcarbamoyloxy)-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid diethyl ester.

To (1S\*, 2S\*, 4S\*, 5R\*, 6R\*)-2-tert-Butoxycarbonylamino-4-hydroxy-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid diethyl ester (0.30g, 0.84 mmol) in 10 mL of toluene, was added in one portion, 2-fluorophenyl isocyanate (0.26g, 1.86 mmol). The reaction was set to reflux for 20 hours. Upon completion, indicated by TLC, the reaction was concentrated under vacuum. Purification of (1S\*, 2S\*, 4S\*, 5R\*, 6R\*)-2-tert-Butoxycarbonylamino-4-(2-fluoro-phenylcarbamoyloxy)-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid diethyl ester was by PC-TLC (10%EtOAc/Hexane to 25% EtOAc/Hexane), with a 94% yield (0.39g, 0.79 mmol).

FDMS: M<sup>+</sup>-1=493.

-127-

B. (1S\*, 2S\*, 4S\*, 5R\*, 6R\*)-2-Amino-4-(2-fluoro-phenylcarbamoyloxy)-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid.

A solution of the product from step (a) (0.36g, 0.73 mmol), in EtOAc was chilled to 0°C. The reaction solution was saturated with HCl (g) and allowed to react until the Boc group was cleaved as indicated by TLC. The reaction was then concentrated under vacuum and rediluted in THF (12mL). 1N NaOH (12mL, 12 mmols) was added and the mixture was allowed to stir overnight (pH=13-14). The reaction mixture was titrated with 1N HCl to a pH=7, and the organics were removed under vacuum. The title compound was isolated by isoelectric point crystalization from H<sub>2</sub>O, pH 3, in a 57% yield (0.14g, 0.41 mmol) (A small amount of IPA was used to transfer the crystals from the flask to drying funnel.).

m.p.=232-233°C.

FDMS: M<sup>+</sup>+1=339.


Anal. Calcd. For C<sub>15</sub>H<sub>15</sub>N<sub>2</sub>O<sub>6</sub> F. 0.5 IPA, 0.6 H<sub>2</sub>O: C, 52.27; H, 5.37; N, 7.39.

Found: C, 52.21; H, 5.15; N, 7.39.



Example 67

Synthesis of (1S\*, 2S\*, 4S\*, 5R\*, 6R\*)-2-Amino-4-(3-fluorophenylcarbamoyloxy)-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid



The title compound was prepared by the two step procedure in Example Bick. (1S\*, 2S\*, 4S\*, 5R\*, 6R\*)-2-tert-Butoxycarbonylamino-4-hydroxy-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid diethyl ester (0.30g, 0.84 mmol) was reacted with 3-fluorophenyl isocyanate (0.19g, 1.35 mmol) in step (a). Purification of the desired intermediate was by PC-TLC (10% EtOAc/Hexane to 25% EtOAc/Hexane) with a 65% yield (0.27g, 0.55 mmol). (1S\*, 2S\*, 4S\*, 5R\*, 6R\*)-2-tert-Butoxycarbonylamino-4-(3-fluorophenylcarbamoyloxy)-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid diethyl ester (0.24g, 0.49 mmol) was converted into the title compound by step (b) using isoelectric point crystalization from H<sub>2</sub>O, pH 3 in a 54% yield (0.09g, 0.27 mmol).

m.p.=239-240°C.


FDMS: M<sup>+</sup>+1=339.

Anal. Calcd. For C<sub>15</sub>H<sub>15</sub>N<sub>2</sub>O<sub>6</sub> F.0.2 H<sub>2</sub>O: C, 52.69; H, 4.54; N, 8.19.

Found: C, 52.49; H, 4.78; N, 8.07.

Example 68

Synthesis of (1S\*, 2S\*, 4S\*, 5R\*, 6R\*)-2-Amino-4-(4-fluorophenylcarbamoyloxy)-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid



The title compound was prepared by the two step procedure in Example Bick. (1S\*, 2S\*, 4S\*, 5R\*, 6R\*)-2-*tert*-Butoxycarbonylamino-4-hydroxy-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid diethyl ester (0.30g, 0.84 mmol) was reacted with 4-fluorophenyl isocyanate (0.18g, 1.11 mmol) in step (a). Purification of the desired intermediate was by PC-TLC (10% EtOAc/Hexane to 25% EtOAc/Hexane) with a 43% yield (0.28g, 0.57 mmol). (1S\*, 2S\*, 4S\*, 5R\*, 6R\*)-2-*tert*-Butoxycarbonylamino-4-(4-fluorophenylcarbamoyloxy)-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid diethyl ester (0.28g, 0.57 mmol) was converted into the title compound by step (b) using isoelectric point crystalization from H<sub>2</sub>O, pH 3 in a 41% yield (0.08g, 0.23 mmol).

m.p.=259-260°C.


FDMS: M<sup>+</sup>+1=339.

Anal. Calcd. For C<sub>15</sub>H<sub>15</sub>N<sub>2</sub>O<sub>6</sub> F. 0.1 NaCl: C, 52.35; H, 4.39; N, 8.14.

Found: C, 51.98; H, 4.01; N, 7.96.

Example 69

Synthesis of (1S\*, 2S\*, 4S\*, 5R\*, 6R\*)-2-Amino-4-(3-chloro-phenylcarbamoyloxy)-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid



The title compound was prepared by the two step procedure in Example Bick. (1S\*, 2S\*, 4S\*, 5R\*, 6R\*)-2-tert-Butoxycarbonylamino-4-hydroxy-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid diethyl ester (0.30g, 0.84 mmol) was reacted with meta-chlorophenyl isocyanate (0.16g, 1.01 mmol) in step (a). Purification of the desired intermediate was by PC-TLC (10% EtOAc/Hexane to 25% EtOAc/Hexane) with a 85% yield (0.37g, 0.72 mmol). (1S\*, 2S\*, 4S\*, 5R\*, 6R\*)-2-tert-Butoxycarbonylamino-4-(3-chloro-phenylcarbamoyloxy)-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid diethyl ester (0.37g, 0.72 mmol) was converted into the title compound by step (b) using isoelectric point crystalization from H<sub>2</sub>O, pH 3 in a 47% yield (0.12g, 0.34 mmol).

m.p.=237-238°C.


FDMS: M<sup>+</sup>+1=355.

Anal. Calcd. For C<sub>15</sub>H<sub>15</sub>N<sub>2</sub>O<sub>6</sub> Cl.0.5 H<sub>2</sub>O: C, 49.53; H, 4.43; N, 7.70.

Found: C, 49.53; H, 4.36; N, 7.59.

Example 70

Synthesis of (1S\*, 2S\*, 4S\*, 5R\*, 6R\*)-2-Amino-4-(4-chloro-phenylcarbamoyloxy)-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid



The title compound was prepared by the two step procedure in Example Bick. (1S\*, 2S\*, 4S\*, 5R\*, 6R\*)-2-*tert*-Butoxycarbonylamino-4-hydroxy-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid diethyl ester (0.30g, 0.84 mmol) was reacted with *p*-chlorophenyl isocyanate (0.16g, 1.01 mmol) in step (a). Purification of the desired intermediate was by PC-TLC (10% EtOAc/Hexane to 25% EtOAc/Hexane) with a 74% yield (0.32g, 0.62 mmol). (1S\*, 2S\*, 4S\*, 5R\*, 6R\*)-2-*tert*-Butoxycarbonylamino-4-(4-chloro-phenylcarbamoyloxy)-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid diethyl ester (0.28g, 0.54 mmol) was converted into the title compound by step (b) using isoelectric point crystalization from H<sub>2</sub>O, pH 3, in a 73% yield (0.14g, 0.39 mmol).

m.p.=258-259°C.


FDMS: M<sup>+</sup>-1=353.

Anal. Calcd. For C<sub>15</sub>H<sub>15</sub>N<sub>2</sub>O<sub>6</sub> Cl.0.2 H<sub>2</sub>O: C, 50.27; H, 4.33; N, 7.82.

Found: C, 50.12; H, 4.25; N, 7.72.

Example 71

Synthesis of (1S\*, 2S\*, 4S\*, 5R\*, 6R\*)-2-Amino-4-(naphthalen-1-ylcarbamoxyloxy)-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid



The title compound was prepared by the two step procedure in Example Bick. (1S\*, 2S\*, 4S\*, 5R\*, 6R\*)-2-tert-Butoxycarbonylamino-4-hydroxy-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid diethyl ester (0.30g, 0.84 mmol) was reacted with 1-naphthyl isocyanate (0.21g, 1.26 mmol) in step (a). Purification of the desired intermediate was by PC-TLC (10% EtOAc/Hexane to 25% EtOAc/Hexane) with a 61% yield. By step (b), using LiOH for the ester hydrolysis, (1S\*, 2S\*, 4S\*, 5R\*, 6R\*)-2-tert-Butoxycarbonylamino-4-(naphthalen-1-ylcarbamoxyloxy)-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid diethyl ester (0.31g, 0.59 mmol) was converted to the Boc-protected di-acid. Purification of the desired intermediate was by PC-TLC (0-2 % AcOH in 50% EtOAc/Hexane) for a 61% yield (0.17g, 0.36 mmol). The Boc group was cleaved, giving the title compound in a 63% yield (0.08g, 0.23 mmol).

m.p.=238-239°C.

FDMS: M<sup>+</sup>+1=371.


-133-

Anal. Calcd. For  $C_{19}H_{18}N_2O_6 \cdot 0.1 H_2O$ : C, 61.32; H, 4.93; N, 7.53.

Found: C, 61.67; H, 4.86; N, 7.29.

### Example 72

Synthesis of (1S\*, 2S\*, 4S\*, 5R\*, 6R\*)-2-Amino-4-(naphthalen-2-ylcarbamoxyloxy)-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid



The title compound was prepared by the two step procedure in Example Bick. (1S\*, 2S\*, 4S\*, 5R\*, 6R\*)-2-tert-Butoxycarbonylamino-4-hydroxy-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid diethyl ester (0.25g, 0.70 mmol) was reacted with 2-naphthyl isocyanate (0.26g, 1.51 mmol) in step (a). Purification of the desired intermediate was by PC-TLC (10% EtOAc/Hexane to 25% EtOAc/Hexane) with a 68% yield (0.25g, 0.47 mmol). By step (b), using LiOH for the ester hydrolysis, (1S\*, 2S\*, 4S\*, 5R\*, 6R\*)-2-tert-Butoxycarbonylamino-4-(naphthalen-2-ylcarbamoxyloxy)-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid diethyl ester (0.28g, 0.53 mmol) was converted to the Boc-protected diacid followed by cleavage of the Boc group. The title

-134-

compound was isolated using isoelectric point crystalization from H<sub>2</sub>O, pH 3, in a 20% yield (0.04g, 0.11 mmol).

m.p.=238-239°C.


FDMS: M<sup>+</sup>-1=369.

Anal. Calcd. For C<sub>19</sub>H<sub>18</sub>N<sub>2</sub>O<sub>6</sub>.0.8 HCl: C, 57.12; H, 4.74; N, 7.01.

Found: C, 57.01; H, 4.82; N, 6.85.

### Example 73

Synthesis of (1S\*,2S\*,4S\*,5R\*,6R\*) 2-Amino-4-(2,4-dichlorophenoxy)bicyclo[3.1.0]hexane-2,6-dicarboxylic acid



A. (1S\*,2S\*,4S\*,5R\*,6R\*) Diethyl 2-(N-t-butyloxycarbonylamino)-4-[(2,4-dichlorophenoxy)bicyclo[3.1.0]hexane-2,6-dicarboxylic acid.

A 0°C solution of (1S\*,2S\*,4S\*,5R\*,6R\*) diethyl 2-(N-t-butyloxycarbonylamino)-4-hydroxybicyclo[3.1.0]hexane-2,6-dicarboxylic acid (1.0 g, 2.8 mmol—synthesis previously described in US 5,958,960 example 8a) in N,N-dimethylformamide (10 mL) was treated in one portion with sodium hydride (0.13 g, 3.4 mmol) and stirred vigorously for 15 minutes. The resulting reaction mixture was treated in one portion with 1,3-dichloro-4-fluorobenzene (0.92 g, 5.6

-135-

mmol) and allowed to warm to room temperature as it stirred overnight. The reaction mixture was partitioned between EtOAc and water and the product extracted with EtOAc. All organic layers were combined, washed with brine, dried over  $\text{MgSO}_4$ , concentrated in vacuo and purified by  $\text{SiO}_2$  chromatography (PC-TLC: 10 % EtOAc/hexanes to 50 % EtOAc/hexanes) to yield 0.63 g (1.3 mmol, 45 %) of a white foam.

FDMS:  $M^+$  = 502, 504.

Anal. calcd. for  $\text{C}_{23}\text{H}_{29}\text{Cl}_2\text{NO}_7 \cdot 0.1 \text{ eq. CH}_2\text{Cl}_2$ : C, 54.31; H, 5.76; N, 2.74.

Found: C, 54.56; H, 5.75; N, 2.93.

B. (1S\*,2S\*,4S\*,5R\*,6R\*) 2-Amino)-4-(2,4-dichlorophenoxy)-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid.

A 0 °C solution of the product of example 20a (0.62 g, 1.23 mmol) in EtOAc (35mL) was purged with anhydrous HCl gas until the solution was saturated. The resulting reaction mixture was stirred at 0 °C for 4 hours and concentrated to dryness in vacuo to afford the HCl salt. The resulting solids were reconstituted in a 1:1 mixture of 1N NaOH:THF (20mL total volume) and stirred at room temperature overnight. The reaction mixture was adjusted to pH = 7 with 1N HCl and solids removed and filtrate concentrated to dryness in vacuo. The solids were reconstituted in minimal amount of water and the product isolated by isoelectric precipitation at pH = 2-3. The product was dried under vacuum at 80 °C to afford 0.38 g (1.1 mmol, 89 %) of the desired product as a white solid.

mp = >250 °C.



-136-


FDMS:  $M^+$  = 346, 348.

Anal. calcd. for  $C_{14}H_{13}Cl_2NO_5 \cdot 0.3 \text{ eq. } H_2O$ : C, 47.83; H, 3.90; N, 3.98.

Found: C, 47.55; H, 3.61; N, 3.97.

#### Example 74

Synthesis of (1S\*,2S\*,4R\*,5R\*,6S\*) 2-Amino-4-phenyl-carbamoylmethylbicyclo[3.1.0]hexane-2,6-dicarboxylic acid



A. 4(Z/E)-(1S\*,2S\*,5R\*,6S\*) Diethyl-2-(*tert*-butoxy carbonylamino)-4-allyloxycarbonylmethylenebicyclo[3.1.0]hexane-2,6-dicarboxylate.

To a 0 °C solution of allyl diethylphosphonoacetate (16.8 mmol, 3.8 g.) in anhydrous THF, was added 1M NaHMDS (16.8 mmol) by dropwise addition. The resulting reaction mixture was stirred at 0 °C for 1 h, after which time a solution of (1S\*,2S\*,5R\*,6R\*) diethyl 2-N-*tert*-butoxycarbonylamino-4-oxobicyclo[3.1.0]hexane-2,6-dicarboxylate (previously disclosed US 5,958,960 example 5a-11.25 mmol, 3.9 g) in THF was added dropwise. The reaction mixture was allowed to room temperature as it stirred overnight. The reaction was quenched with saturated aqueous  $NH_4Cl$  solution and the product extracted with ethyl acetate. The combined organic extracts were dried over sodium sulfate, concentrated to dryness in vacuo, and purified by  $SiO_2$  chromatography. (50 %

-137-

EtOAc/ hexanes) to afford 4.8 g (10.9 mmol, 97%) of the title compound as a (1:1.5) mixture of diastereoisomers as colorless oil.

$^1\text{H-NMR}$  ( $\text{CDCl}_3$ )  $\delta$ : 5.91-5.77 (m, 4H), 5.45-5.14 (m, 6H), 4.60-4.55 (m, 4H), 4.25-4.00 (m, 8H), 3.89 (d, 1H,  $J=19.8$  Hz), 3.64 (dd, 1H,  $J=2,7$  and 5.9 Hz), 3.25 (d, 1H  $J=17.4$  Hz), 2.79 (m, 1H), 2.63-2.57 (m, 2H), 2.24-2.08 (m, 2H), 2.07-2.02 (m, 2H), 1.4 (s, 9 H), 1.39 (s, 9H), 1.22 (dt, 12H) ppm.  $^{13}\text{C-NMR}$  ( $\text{CDCl}_3$ )  $\delta$ : 172.2, 172.0, 170.3, 170.1, 165.6, 165.3, 160.1, 159.9, 132.2, 117.7, 113.5, 113.0, 65.3, 64.5, 63.5, 61.8, 61.7, 61.1, 60.2, 41.3, 38.5, 36.6, 35.8, 35.3, 31.5, 28.0, 25.4, 25.1, 14.0, 13.9 ppm. MS(ES) for  $\text{C}_{22}\text{H}_{31}\text{NO}_8$   $[\text{M}+\text{H}]^+$ : 438.18;  $[\text{M}-\text{H}]^-$ : 436.4.

B. 4(Z/E)-(1S\*,2S\*,5R\*,6S\*) Diethyl-2-(tert-butyloxycarbonylamino)-4-carboxymethylene bicyclo[3.1.0] hexane-2,6-dicarboxylate.

Method B1) An EtOH/ $\text{H}_2\text{O}$  (9:1) solution of the mixture of compounds from Example 74A (4.3 mmol, 1.9 g) was treated with Wilkinson's catalyst (Chlorotris triphenylphosphine)rhodium (I) (0.4 mmol, 0.370 g) and the resulting reaction mixture warmed at reflux for 5 hours. Upon cooling to room temperature, the mixture reaction was filtered through a pad of celite to remove the catalyst. The solvent was removed in vacuo and the crude reaction was passed through a short pad of  $\text{SiO}_2$  to afford the title compound as a (1:1.5) mixture of isomers (1.28 g, 3.2 mmol, 73%).

-138-

Method B2) Tetrakis(triphenylphosphine) Palladium (0) (0.150 g, 0.13 mmol) and pyrrolidine (0.48 g, 6.7 mmol) were sequentially added to a solution of the mixture of isomers from Example 74A (4.5 mmol, 2g) in  $\text{CH}_2\text{Cl}_2$  and the resulting reaction mixture warmed at reflux over night. The reaction was allowed to cool to room temperature and washed with  $\text{NaHSO}_4$  and brine. The organic layer was concentrated to dryness, the residue diluted in MeOH, and the remaining insolubles removed via vacuum filtration. The filtrate was concentrated in vacuo and purified by filtration through a short pad of  $\text{SiO}_2$  to afford a 1:1.5 mixture of the free carboxylic acids (0.9 g, 2.2 mmol, 53 %).

$^1\text{H-NMR}$  ( $\text{CDCl}_3$ )  $\delta$ : 5.92 (s, 1H), 5.78 (s, 1H), 5.33 (brs, 2H), 4.25-4.09 (m, 8H), 3.95 d, 1H,  $J = 19.0$  Hz), 3.64 (dd, 1H,  $J = 2.6$  and  $6.1$  Hz), 2.77 (dd, 1H,  $J = 3.2$  and  $5.9$  Hz), 2.68-2.64 (m, 1H), 2.27-2.00 ((m, 4H), 1.43 (s, 18 H), 1.31-1.21 (m, 12H) ppm.

C. 4(Z/E), (1S\*,2S\*,5R\*,6S\*) Diethyl 2-tert-butyloxy-carbonylamino-4-phenyl carbamoyl methylenebicyclo[3.1.0]-hexane-2,6-dicarboxylate.

A room temperature solution of the product of Example 74B (0.5 g, 1.25 mmol) in  $\text{CH}_2\text{Cl}_2$  was treated consecutively with oxalyl chloride (0.239 g, 1.89 mmol) and a catalytic amount of DMF. After the reaction mixture was stirred for one hour, the solvent was removed in vacuo. The crude product was reconstituted in  $\text{CH}_2\text{Cl}_2$  and then a solution of aniline (0.232 g, 2.5 mmol) was added dropwise. The reaction mixture was allowed to stir at room temperature until the reaction was judged complete by TLC. The resulting mixture was concentrated to dryness and the crude product purified by

-139-

filtration through a short pad of SiO<sub>2</sub> to afford the title compounds (0.422 g, 0.89mmol) 71 % as a mixture of diastereoisomers which were used in the next step without further separation.

MS(ES) for C<sub>25</sub>H<sub>32</sub>N<sub>2</sub>O<sub>7</sub> [M+H]<sup>+</sup>: 473.2

D. (1S\*,2S\*,4R\*,5R\*,6S\*) Diethyl 2-*tert*-butyloxycarbonyl-amino-4-phenylcarbamoylmethylbicyclo[3.1.0]hexane-2,6-dicarboxylate.

A room temperature solution of the product of Example 74C (0.422 g, 0.89mmol) and 10% Pd/C (0.021 g, 5 wt %) in 95% EtOH was vigorously stirred under an hydrogen atmosphere (1 atm) overnight. The catalyst was removed via vacuum filtration through celite and the filtrate concentrated in vacuo to afford the title compound (0.34 g, 0.45 mmol, 80%).

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 8.03 (s, 1H), 7.54 (d, 2H, J= 7.8 Hz), 7.30 (t, 2H), 7.07 (t, 1H, J= 7.5 Hz), 5.22 (s, 1H), 4.25-4.20 (m, 2H), 4.12 (q, 2H, J= 7.2 Hz), 2.98-2.86 (m, 1H), 2.67 (d, 1H, J= 15 Hz), 2.54 (d, 2H, J= 8.3 Hz), 2.36 (dd, 1H, J= 2.7 and 5.9 Hz), 1.88 (dd, 1H, J= 3.5 and 6.0 Hz), 1.77 (t, 1H, J= 2.9 Hz), 1.52-1.20 (m, 15 H) ppm.

MS(ES) for C<sub>25</sub>H<sub>34</sub>N<sub>2</sub>O<sub>7</sub> [M+H]<sup>+</sup>: 475.07; [M-H]<sup>-</sup>: 473.23.

E. (1S\*,2S\*,4R\*,5R\*,6S\*) Diethyl 2-amino-4-phenylcarbamoylmethylbicyclo[3.1.0] hexane-2,6-dicarboxylate.

The product of example 74D (0.297 g, 0.62 mmol) was treated with a saturated solution of EtOAc/HCl and stirred at room temperature over night. The reaction mixture was adjusted to

-140-

pH = 7-8 with  $\text{NaHCO}_3$ , and the product extracted with EtOAc. The crude product was purified by flash chromatography to afford the title compound (0.152 g, 0.4 mmol, 65%).

$^1\text{H}$ -NMR ( $\text{CDCl}_3$ )  $\delta$ : 8.18 (s, 1H, NHCO), 7.55 (d, 2H,  $J$  = 12 Hz), 7.24-7.12 (m, 2H), 6.89-6.78 (m, 1H), 4.27 (q, 2H,  $J$  = 7.1 Hz), 4.11 (q, 2H,  $J$  = 7.0 Hz), 2.84-2.75 (m, 1H), 2.52-2.25 (m, 4H), 1.86-1.84 (m, 3H), 1.77 (t, 1H,  $J$  = 2.9 Hz), 1.40-1.21 (dt, 6H) ppm.

MS(ES) for  $\text{C}_{20}\text{H}_{26}\text{N}_2\text{O}_5$   $[\text{M}+\text{H}]^+$ : 375.04;  $[\text{M}-\text{H}]^-$ : 373.15

F. (1S\*,2S\*,4R\*,5R\*,6S\*)- 2-amino-4-phenylcarbamoylmethyl-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid.

A solution of the product of Example 74E (0.083 g, 0.22 mmol) in THF was treated with 1N NaOH ( 0.55 mmol) and stirred at room temperature over night. The solvent was removed in vacuo and the crude product reconstituted in distilled water and acidified with 1N HCl. The product was isolated by isoelectric precipitation at pH = 3-5. The product was collected via vacuum filtration, washed with water and air dried to afford the title compound (0.035 g, 0.11mmol, 51%).

$^1\text{H}$ - NMR ( $\text{D}_2\text{O}$  + KOD)  $\delta$ : 7.37-7.34 (m, 4H), 7.22-7.17 (m, 1H), 2.64-2.37 (m, 3H), 2.03 (dd, 1H,  $J$  = 3.3 and 6.3 Hz), 1.88 (d, 1H,  $J$  = 14.9 Hz), 1.67 (dd, 1H,  $J$  = 2.5 and 5.8 Hz), 1.46-1.33 (m, 2H) ppm.


$^{13}\text{C}$ - NMR ( $\text{D}_2\text{O}$  + KOD )  $\delta$ : 183.7, 176.2, 138.7, 131.0, 127.4, 124.0, 67.9, 43.6, 42.7, 39.7, 37.3, 33.6, 26.8 ppm.

MS(ES) for  $\text{C}_{16}\text{H}_{18}\text{N}_2\text{O}_5$   $[\text{M}-\text{H}]^-$ : 317.22

Pf > 190° dec.

Example 75

Synthesis of (1S\*,2S\*,4R\*,5R\*,6S\*) 2-Amino-4-(o-methoxy)-phenylcarbamoylmethyl bicyclo[3.1.0]hexane-2,6-dicarboxylic acid



A. 4(Z/E), (1S\*,2S\*,5R\*,6S\*) Diethyl 2-*tert*-butoxy-carbonylamino-4-(o-methoxy)phenylcarbamoylmethylene bicyclo[3.1.0]hexane-2,6-dicarboxylate.

A room temperature solution of the product of Example 74B (0.5 g, 1.25 mmol) in CH<sub>2</sub>Cl<sub>2</sub> was treated consecutively with oxalyl chloride (0.239 g, 1.89 mmol) and a catalytic amount of DMF. After the reaction mixture was stirred for one hour, the solvent was removed in vacuo. The crude product was reconstituted in CH<sub>2</sub>Cl<sub>2</sub> and then a solution of o-anisidine (0.307 g, 2.5 mmol) was added dropwise. The reaction mixture was allowed to stir at room temperature until the reaction was judged complete by TLC. The resulting mixture was concentrated to dryness and the crude product purified by filtration through a short pad of SiO<sub>2</sub> to afford the title compounds (0.260 g, 0.51 mmol) 40 % as a mixture of diastereoisomers which were used in the next step without further separation.

MS(ES) for C<sub>26</sub>H<sub>34</sub>N<sub>2</sub>O<sub>8</sub> [M+H]<sup>+</sup>: 502.11

-142-

B. (1S\*,2S\*,4R\*,5R\*,6S\*) Diethyl 2-*tert*-butyloxycarbonyl-amino-4-(*o*-methoxy) phenylcarbamoylmethylbicyclo[3.1.0]-hexane-2,6-dicarboxylate.

A room temperature solution of the product of Example 75A (0.260 g, 0.51mmol) and 10% Pd/C (0.021 g, 5 wt %) in 95% EtOH was vigorously stirred under an hydrogen atmosphere (1 atm) over night. The catalyst was removed via vacuum filtration through celite and the filtrate concentrated in vacuo to afford the title compound (0.150 g, 0.29 mmol, 60%).

C<sub>26</sub>H<sub>36</sub>N<sub>2</sub>O<sub>8</sub> [M+H]<sup>+</sup>: 505.09. [M-H]<sup>-</sup>: 503.28

C. (1S\*,2S\*,4R\*,5R\*,6S\*) Diethyl 2-Amino-4-(*o*-methoxy) phenylcarbamoylmethyl bicyclo[3.1.0]hexane-2,6-dicarboxylate.

The title compound was prepared as example 74E to give the product in 57% yield.

<sup>1</sup>H NMR (CDCl<sub>3</sub>) δ: 8.32 (d, 1H, J= 7.6Hz), 7.97 (s, 1H), 7.02-6.82 (m, 3H), 4.25 (q, 2H, J= 7.1 Hz), 4.10 (dq, 2H, J= 1.3 and 7.2 Hz), 3.85 (s, 3H), 2.87-2.82 (m, 1H), 2.51-2.46 (m, 2H), 2.32 (dd, 1H, J= 2.9 and 6.0 Hz), 2.21 (d 1H, J= 14.8 Hz), 1.91 (dd, 1H, J= 3.0 and 6.1 Hz), 1.76-1.72 (m, 3H), 1.44 (dd, 1H, J= 8.3 and 14.8 Hz), 1.37-1.20 (m, 6H) ppm.

MS(ES) for C<sub>21</sub>H<sub>28</sub>N<sub>2</sub>O<sub>6</sub> [M+H]<sup>+</sup>: 405.04; [M-H]<sup>-</sup>: 403.16

D. (1S\*,2S\*,4R\*,5R\*,6S\*) 2-Amino-4-(*o*-methoxy)phenyl carbamoylmethyl bicyclo[3.1.0]hexane-2,6-dicarboxylic acid.

-143-

A solution of the product of Example 75C (0.068 g, 0.16 mmol) in THF was treated with 1N NaOH (0.33 mmol) and stirred at room temperature over night. The solvent was removed in vacuo and the crude product reconstituted in distilled water and acidified with 1N HCl. The product was isolated by isoelectric precipitation at pH = 3-5. The product was collected via vacuum filtration, washed with water and air dried to afford the title compound (0.030 g, 0.08mmol, 52%).


$^1\text{H}$ - NMR (MeOD+Pyr)  $\delta$ : 7.36 (dd, 1H,  $J$  = 1.4 and 7.7 Hz); 7.20 (m, 1H); 7.03-6.88 (m, 2H); 3.76 (s, 3H); 2.57-2.35 (m, 3H), 2.01 (dd, 1H,  $J$  = 2.8 and 5.8 Hz), 1.89 (d, 1H,  $J$  = 14.4 Hz); 1.72 (dd, 1H,  $J$  = 2.9 and 6.0 Hz); 1.47-1.36 (m, 2H) ppm.

$^{13}\text{C}$ - NMR ( $\text{D}_2\text{O}$ +KOD)  $\delta$ : 183.4, 181.9, 174.8, 152.3, 127.4, 125.8, 121.6, 112.3, 66.0, 55.9, 50.4, 41.1, 38.0, 35.6, 31.7, 25.1 ppm

MS(ES) for  $\text{C}_{17}\text{H}_{20}\text{N}_2\text{O}_6$   $[\text{M}+\text{H}]^+$ : 348.99  $[\text{M}-\text{H}]^-$ : 347.08

### Example 76

Synthesis of (1S\*,2S\*,4R\*,5R\*,6S\*) 2-Amino-4-(o-methyl) phenylcarbamoylmethyl bicyclo[3.1.0]hexane-2,6-dicarboxylic acid





-144-

A. 4(Z/E), (1S\*,2S\*,5R\*,6S\*) Diethyl 2-*tert*-butyloxy carbonylamino-4-(*o*-methyl)phenylcarbamoyle methylene bicyclo[3.1.0]hexane-2,6-dicarboxylate.

A room temperature solution of the product of Example 74B (0.5 g, 1.25 mmol) in CH<sub>2</sub>Cl<sub>2</sub> was treated consecutively with oxalyl chloride (0.239 g, 1.89 mmol) and a catalytic amount of DMF. After the reaction mixture was stirred for one hour, the solvent was removed in vacuo. The crude product was reconstituted in CH<sub>2</sub>Cl<sub>2</sub> and then a solution of *o*-toluididine (0.268 g, 2.5 mmol) was added dropwise. The reaction mixture was allowed to stir at room temperature until the reaction was judged complete by TLC. The resulting mixture was concentrated to dryness and the crude product purified by filtration through a short pad of SiO<sub>2</sub> to afford the title compounds (0.370 g, 0.79 mmol) 60 % as a mixture of diastereoisomers which were used in the next step without further separation.

B. (1S\*,2S\*,4R\*,5R\*,6S\*) Diethyl 2-*tert*-butyloxy carbonylamino-4-(*o*-methyl)phenylcarbamoylemethyl bicyclo[3.1.0]hexane-2,6-dicarboxylate.

A room temperature solution of the product of Example 76A (0.370 g, 0.76 mmol) and 10% Pd/C (0.20 g, 5 wt %) in 95% EtOH was vigorously stirred under an hydrogen atmosphere (1 atm) over night. The catalyst was removed via vacuum filtration through celite and the filtrate concentrated in vacuo to afford the title compound (0.224 g, 0.45 mmol, 62%).

C. (1S\*,2S\*,4R\*,5R\*,6S\*) Diethyl 2-Amino-4-(*o*-methyl)phenylcarbamoylemethylbicyclo[3.1.0]hexane-2,6-dicarboxylate.

-145-

The title compound was prepared using the same procedure as in Example 74E to give the product.

$^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$ : 7.83 (d, 1H,  $J = 7.7\text{Hz}$ ), 7.71 (s, 1H), 7.20-6.99 (m, 3H), 4.26-4.04 (dq, 4H), 2.87-2.75 (m, 1H), 2.50-2.21 (m, 8H), 1.86 (dd, 1H,  $J = 3.0$  and  $6.1\text{Hz}$ ), 1.76 (br s, 3H), 1.41-1.18 (m, 6H) ppm.

MS(ES) for  $\text{C}_{21}\text{H}_{28}\text{N}_2\text{O}_5$   $[\text{M}+\text{H}]^+$ : 389.06;  $[\text{M}-\text{H}]^-$ : 387.16


D. (1S\*,2S\*,4R\*,5R\*,6S\*) 2-Amino-4-(o-methyl)phenyl carbamoylmethyl bicyclo[3.1.0] hexane-2,6-dicarboxylic acid.

A solution of the product of Example 76C (0.090 g, 0.23 mmol) in THF was treated with 1N NaOH (0.46 mmol) and stirred at room temperature over night. The solvent was removed in vacuo and the crude product reconstituted in distilled water and acidified with 1N HCl. The product was isolated by isoelectric precipitation at pH = 3-5. The product was collected via vacuum filtration, washed with water and air dried to afford the title compound (0.030 g, 0.09mmol, 39 %).  $^1\text{H}$ - NMR ( $\text{MeOD}+\text{Pyr}$ )  $\delta$ : 7.26-7.15 (m, 4H); 2.62-2.42 (m, 3H), 2.15 (s, 3H); 2.05 (dd, 1H,  $J = 3.1$  and  $6.1\text{Hz}$ ), 1.94 (d, 1H,  $J = 14.5\text{Hz}$ ); 1.72 (dd, 1H,  $J = 2.6$  and  $6.0\text{Hz}$ ); 1.51-1.40 (m, 2H) ppm.  $^{13}\text{C}$ - NMR ( $\text{D}_2\text{O}+\text{KOD}$ )  $\delta$ : 183.4, 181.9, 174.9, 134.7, 134.5, 130.7, 127.5, 126.9, 126.5, 66.1, 41.1, 37.9, 35.5, 31.9, 25.0, 17.0 ppm

MS(ES) for  $\text{C}_{17}\text{H}_{20}\text{N}_2\text{O}_5$   $[\text{M}+\text{H}]^+$ : 332.99  $[\text{M}-\text{H}]^-$ : 331.09

Example 77

Synthesis of (1S\*,2S\*,4R\*,5R\*,6S\*) 2-Amino-4-(o-trifluoromethyl)phenylcarbamoylmethyl bicyclo[3.1.0]hexane-2,6-dicarboxylic acid



A. 4(Z/E), (1S\*,2S\*,5R\*,6S\*) Diethyl 2-*tert*-butoxy-carbonylamino-4-(o-trifluoromethyl)phenyl carbamoylmethylene bicyclo[3.1.0]hexane-2,6-dicarboxylate.

A room temperature solution of the product of Example 74B (0.5 g, 1.25 mmol) in CH<sub>2</sub>Cl<sub>2</sub> was treated consecutively with oxalyl chloride (0.239 g, 1.89 mmol) and a catalytic amount of DMF. After the reaction mixture was stirred for one hour, the solvent was removed in vacuo. The crude product was reconstituted in CH<sub>2</sub>Cl<sub>2</sub> and then a solution of *o*-trifluoromethyl aniline (0.402 g, 2.5 mmol) was added dropwise. The reaction mixture was allowed to stir at room temperature until the reaction was judged complete by TLC. The resulting mixture was concentrated to dryness and the crude product purified by filtration through a short pad of SiO<sub>2</sub> to afford the title compounds (0.480 g, 0.88 mmol) 79 % as a mixture of diastereoisomers which were used in the next step without further separation. MS(ES) for C<sub>26</sub>H<sub>31</sub>F<sub>3</sub>N<sub>2</sub>O<sub>7</sub> [M+H]<sup>+</sup>: 541.21 [M-H]<sup>-</sup>: 539.24.

-147-

B. (1S\*,2S\*,4R\*,5R\*,6S\*) Diethyl 2-tert-butyloxy-carbonylamino-4-(o-trifluoromethyl)phenyl carbamoylmethyl-bicyclo[3.1.0]hexane-2,6-dicarboxylate.

A room temperature solution of the product of Example 77A (0.370 g, 0.76 mmol) and 10% Pd/C (0.20 g, 5 wt %) in 95% EtOH was vigorously stirred under an hydrogen atmosphere (1 atm) over night. The catalyst was removed via vacuum filtration through celite and the filtrate concentrated in vacuo to afford the title compound

MS(ES) for  $C_{26}H_{33}F_3N_2O_7$   $[M+H]^+$ : 543.09  $[M-H]^-$ : 541.24

C. (1S\*,2S\*,4R\*,5R\*,6S\*) Diethyl 2-Amino-4-(o-trifluoromethyl)phenylcarbamoyl methylbicyclo[3.1.0]hexane-2,6-dicarboxylate.

The title compound was prepared using the same procedure as in Example 74E to give the product.

D. (1S\*,2S\*,4R\*,5R\*,6S\*) 2-Amino-4-(o-trifluoromethyl)phenylcarbamoylmethylbicyclo[3.1.0]hexane-2,6-dicarboxylic acid.

A solution of the product of Example 77C (0.090 g, 0.20 mmol) in THF was treated with 1N NaOH (0.40 mmol) and stirred at room temperature over night. The solvent was removed in vacuo and the crude product reconstituted in distilled water and acidified with 1N HCl. The product was isolated by isoelectric precipitation at pH = 3-5. The product was collected via vacuum filtration, washed with water and air dried to afford the title compound (0.045 g, 0.11mmol, 57 %).


-148-

$^1\text{H}$ - NMR (MeOD+Pyr)  $\delta$ : 7.65 (d, 1H,  $J$  = 7.6 Hz); 7.52 (t, 1H,  $J$  = 7.1 Hz); 7.27 (t, 1H,  $J$  = 7.7 Hz); 7.14 (d, 1H,  $J$  = 7.8 Hz); 2.64-2.34 (m, 3H), 2.04 (dd, 1H,  $J$  = 2.8 and 6.0 Hz), 1.94 (d, 1H,  $J$  = 14.4 Hz); 1.76 (dd, 1H,  $J$  = 2.9 and 6.1 Hz); 1.50-1.39 (m, 2H) ppm.

$^{13}\text{C}$ - NMR ( $\text{D}_2\text{O}+\text{KOD}$ )  $\delta$ : 183.5, 182.1, 174.9, 132.8, 128.8, 126.3, 125.3, 66.1, 42.2, 41.8, 38.4, 35.6, 32.2, 25.2 ppm

### Example 78

Synthesis of (1S\*,2S\*,4R\*,5R\*,6S\*) 2-Amino-4-(o-chloro)phenylcarbamoylmethyl bicyclo[3.1.0]hexane-2,6-dicarboxylic acid



A. 4(Z/E), (1S\*,2S\*,5R\*,6S\*) Diethyl 2-tert-butyloxy-carbonylamino-4-(o-chloro) phenylcarbamoylmethylene bicyclo[3.1.0]hexane-2,6-dicarboxylate.

A room temperature solution of the product of Example 74B (0.5 g, 1.25 mmol) in  $\text{CH}_2\text{Cl}_2$  was treated consecutively with oxalyl chloride (0.239 g, 1.89 mmol) and a catalytic amount of DMF. After the reaction mixture was stirred for one hour, the solvent was removed in vacuo. The crude product was reconstituted in  $\text{CH}_2\text{Cl}_2$  and then a solution of o-chloroaniline (0.318 g, 2.5 mmol) was added dropwise. The reaction mixture was allowed to stir at room temperature

-149-

until the reaction was judged complete by TLC. The resulting mixture was concentrated to dryness and the crude product purified by filtration through a short pad of SiO<sub>2</sub> to afford the title compounds (0.283 g, 0.55 mmol) 44 % as a mixture of diastereoisomers which were used in the next step without further separation.

B. (1S\*,2S\*,4R\*,5R\*,6S\*) Diethyl 2-tert-butyloxy carbonylamino-4-(o-chloro)phenylcarbamoyl methylbicyclo[3.1.0]hexane-2,6-dicarboxylate.

A room temperature solution of the product of Example 78A (0.283 g, 0.55 mmol) and 10% Pd/C (0.014 g, 5 wt %) in 95% EtOH was vigorously stirred under a hydrogen atmosphere (1 atm) over night. The catalyst was removed via vacuum filtration through celite and the filtrate concentrated in vacuo to afford the title compound (0.210 g, 0.41 mmol, 74 %). for C<sub>25</sub>H<sub>33</sub>ClN<sub>2</sub>O<sub>7</sub> [M+H]<sup>+</sup>-Cl: 475.08 [M-H]<sup>-</sup>-Cl: 473.22

C. (1S\*,2S\*,4R\*,5R\*,6S\*) Diethyl 2-Amino-4-(o-chloro) phenylcarbamoyl methylbicyclo[3.1.0]hexane-2,6-dicarboxylate.

The title compound was prepared as in Example 74E to give the product in 67 % yield

<sup>1</sup>H NMR (CDCl<sub>3</sub>)δ: 8.07 (s, 1H), 7.53 (d, 2H, J = 7.9 Hz), 7.33-7.25 (m, 1H), 7.07 (t, 1H, J = 7.3 Hz), 4.26 (q, 2H, J = 7.1 Hz), 4.19-4.06 (m, 2H), 2.87-2.75 (m, 1H), 2.45-2.24 (m, 5H), 1.87-1.75 (m, 4H), 1.34-1.20 (m, 6H) ppm.

MS(ES) for C<sub>20</sub>H<sub>25</sub>ClN<sub>2</sub>O<sub>6</sub> [M+H]<sup>+</sup>-Cl: 375.04; [M-H]<sup>-</sup>-Cl: 373.11

-150-


D. (1S\*,2S\*,4R\*,5R\*,6S\*) 2-Amino-4-(o-chloro)phenyl carbamoylmethyl bicyclo[3.1.0]hexane-2,6-dicarboxylic acid.

A solution of the product of Example 78C (0.150 g, 0.36 mmol) in THF was treated with 1N NaOH ( 0.73 mmol) and stirred at room temperature over night. The solvent was removed in vacuo and the crude product reconstituted in distilled water and acidified with 1N HCl. The product was isolated by isoelectric precipitation at pH = 3-5. The product was collected via vacuum filtration, washed with water and air dried to afford the title compound (0.052 g, 0.14mmol, 40 %).

$^1\text{H}$  NMR ( $\text{CDCl}_3$ ) $\delta$ : 8.07 (s, 1H), 7.53 (d, 2H, J = 7.9 Hz), 7.33-7.25 (m, 1H), 7.07 (t, 1H, J = 7.3 Hz), 4.26 (q, 2H, J= 7.1 Hz ), 4.19-4.06 (m, 2H), 2.87-2.75 (m, 1H), 2.45-2.24 (m, 5H), 1.87-1.75 (m, 4H), 1.34-1.20 (m, 6H) ppm. MS (ES) for  $\text{C}_{20}\text{H}_{25}\text{ClN}_2\text{O}_6$   $[\text{M}+\text{H}]^+/-\text{Cl}$ : 375.04;  $[\text{M}-\text{H}]^-/-\text{Cl}$ : 373.11

Example 79

Synthesis of (1S\*,2S\*,4R\*,5R\*,6S\*) 2-Amino-4-(*m*-methoxy)phenylcarbamoylmethyl bicyclo[3.1.0]hexane-2,6-dicarboxylic acid



A. 4(Z/E), (1S\*,2S\*,5R\*,6S\*) Diethyl 2-*tert*-butyloxy-carbonylamino-4-(*m*-methoxy)phenylcarbamoylmethylene bicyclo[3.1.0]hexane-2,6-dicarboxylate.

A room temperature solution of the product of Example 74B (0.5 g, 1.25 mmol) in CH<sub>2</sub>Cl<sub>2</sub> was treated consecutively with oxalyl chloride (0.239 g, 1.89 mmol) and a catalytic amount of DMF. After the reaction mixture was stirred for one hour, the solvent was removed in vacuo. The crude product was reconstituted in CH<sub>2</sub>Cl<sub>2</sub> and then a solution of *m*-anisidine (0.307 g, 2.5 mmol) was added dropwise. The reaction mixture was allowed to stir at room temperature until the reaction was judged complete by TLC. The resulting mixture was concentrated to dryness and the crude product purified by filtration through a short pad of SiO<sub>2</sub> to afford the title compounds (0.505 g, 1.0 mmol) 80 % as a mixture of diastereoisomers which were used in the next step without further separation.



B. (1S\*,2S\*,4R\*,5R\*,6S\*) Diethyl 2-*tert*-butoxy carbonylamino-4-(*m*-methoxy)phenylcarbamoylmethyl bicyclo[3.1.0]hexane-2,6-dicarboxylate.

A room temperature solution of the product of Example 79A (0.505 g, 1.0 mmol) and 10% Pd/C (0.025 g, 5 wt %) in 95% EtOH was vigorously stirred under an hydrogen atmosphere (1 atm) over night. The catalyst was removed via vacuum filtration through celite and the filtrate concentrated in vacuo to afford the title compound (0.354 g, 0.70 mmol, 70 %).

C. (1S\*,2S\*,4R\*,5R\*,6S\*) Diethyl 2-Amino-4-(*m*-methoxy) phenylcarbamoyl methylbicyclo[3.1.0]hexane-2,6-dicarboxylate.

The title compound was prepared as in Example 74E to give the product

$^1\text{H}$  NMR ( $\text{CDCl}_3$ ) $\delta$ : 8.22 (s, 1H, NHCO), 7.33-7.12 (m, 2H), 6.98 (brd, 2H), 6.61 (dd, 1H,  $J = 2.2$  and 8.0 Hz), 4.23 (q, 2H,  $J = 7.1$  Hz), 4.08 (q, 2H,  $J = 7.1$  Hz), 3.76 (s, 3H), 2.81-2.72 (m, 1H), 2.50-2.21 (m, 5H), 1.87-1.74 (m, 4H), 1.39-1.20 (dt, 6H) ppm.

MS(ES) for  $\text{C}_{21}\text{H}_{28}\text{N}_2\text{O}_6$   $[\text{M}+\text{H}]^+$ : 405.05;  $[\text{M}-\text{H}]^-$ : 403.16

D. (1S\*,2S\*,4R\*,5R\*,6S\*) 2-Amino-4-(*m*-methoxy) phenylcarbamoylmethylbicyclo[3.1.0]hexane-2,6-dicarboxylic acid.

A solution of the product of Example 79C (0.140 g, 0.34 mmol) in THF was treated with 1N NaOH (0.69 mmol) and stirred at room temperature over night. The

-153-

solvent was removed in vacuo and the crude product reconstituted in distilled water and acidified with 1N HCl. The product was isolated by isoelectric precipitation at pH = 3-5. The product was collected via vacuum filtration, washed with water and air dried to afford the title compound (0.085 g, 0.24 mmol, 70 %).

$^1\text{H}$ -NMR ( $\text{D}_2\text{O}+\text{Pyr}$ )  $\delta$ : 6.93 (t, 1H,  $J = 7.9$  and  $8.1$  Hz); 6.75 (s, 1H), 6.67 (d, 1H,  $J = 7.9$  Hz); 6.38 (d, 1H,  $J = 8.1$  Hz); 3.43 (s, 3H); 2.48-2.31 (m, 3H), 1.99 (dd, 1H,  $J = 3.4$  and  $3.5$  Hz), 1.80-1.57 (m, 3H), 1.42 (t, 1H,  $J = 3.0$  Hz) ppm.


$^{13}\text{C}$ -NMR ( $\text{D}_2\text{O}+\text{Pyr}$ )  $\delta$ : 179.8, 176.4, 173.4, 159.3, 138.6, 130.2, 113.8, 110.8, 106.8, 67.8, 55.4, 41.6, 38.4, 38.2, 33.5, 33.3, 26.0 ppm.

MS(ES) for  $\text{C}_{17}\text{H}_{20}\text{N}_2\text{O}_6$   $[\text{M}+\text{H}]^+$ : 348.99  $[\text{M}-\text{H}]^-$ : 347.08

Pf > 200° dec.

Example 80

Synthesis of (1S\*,2S\*,4R\*,5R\*,6S\*) 2-Amino-4-(*m*-methyl)phenylcarbamoylmethyl bicyclo[3.1.0]hexane-2,6-dicarboxylic acid



A. 4(Z/E) (1S\*,2S\*,5R\*,6S\*) Diethyl 2-*tert*-butyloxycarbonyl amino-4-(*m*-methyl)phenylcarbamoylmethylene bicyclo[3.1.0]hexane-2,6-dicarboxylate.

A room temperature solution of the product of Example 74B (0.5 g, 1.25 mmol) in CH<sub>2</sub>Cl<sub>2</sub> was treated consecutively with oxalyl chloride (0.239 g, 1.89 mmol) and a catalytic amount of DMF. After the reaction mixture was stirred for one hour, the solvent was removed in vacuo. The crude product was reconstituted in CH<sub>2</sub>Cl<sub>2</sub> and then a solution of *m*-toluidine (0.267 g, 2.5 mmol) was added dropwise. The reaction mixture was allowed to stir at room temperature until the reaction was judged complete by TLC. The resulting mixture was concentrated to dryness and the crude product purified by filtration through a short pad of SiO<sub>2</sub> to afford the title compounds (0.399 g, 0.81 mmol) 65 % as a mixture of diastereoisomers which were used in the next step without further separation.

B. (1S\*,2S\*,4R\*,5R\*,6S\*) Diethyl 2-*tert*-butoxy carbonylamino-4-(*m*-methyl)phenylcarbamoylmethyl bicyclo[3.1.0]hexane-2,6-dicarboxylate.

A room temperature solution of the product of Example 80A (0.309 g, 0.81 mmol) and 10% Pd/C (0.015 g, 5 wt %) in 95% EtOH was vigorously stirred under an hydrogen atmosphere (1 atm) over night. The catalyst was removed via vacuum filtration through celite and the filtrate concentrated in vacuo to afford the title compound (0.217 g, 0.44 mmol, 70 %).

C. (1S\*,2S\*,4R\*,5R\*,6S\*) Diethyl 2-Amino-4-(*m*-methyl) phenylcarbamoyl methylbicyclo[3.1.0]hexane-2,6-dicarboxylate.

The title compound was prepared as in Example 74E to give the product in 64 % yield

$^1\text{H}$  NMR ( $\text{CDCl}_3$ ) $\delta$ : 8.30(s, 1H, NHCO), 7.50-7.24 (m, 3H), 7.99 (d, 1H,  $J$  = 7.4 Hz), 4.36 (q, 2H,  $J$  = 7.1 Hz), 4.26-4.15 (m, 2H), 2.93-2.85 (m, 1H), 2.62-2.34 (m, 8H), 1.99-1.95 (m, 3H), 1.87 (t, 1H,  $J$  = 3.0 Hz), 1.51-1.29 (dt, 6H) ppm.  
MS(ES) for  $\text{C}_{21}\text{H}_{28}\text{N}_2\text{O}_5$   $[\text{M}+\text{H}]^+$ : 389.05;  $[\text{M}-\text{H}]^-$ : 387.17

D. (1S\*,2S\*,4R\*,5R\*,6S\*) 2-Amino-4-(*m*-methyl) phenylcarbamoylmethylbicyclo [3.1.0]hexane-2,6-dicarboxylic acid.

A solution of the product of Example 80C (0.100 g, 0.25 mmol) in THF was treated with 1N NaOH (0.51 mmol) and stirred at room temperature over night. The solvent was removed in vacuo and the crude product reconstituted in distilled water and acidified with 1N

-156-

HCl. The product was isolated by isoelectric precipitation at pH = 3-5. The product was collected via vacuum filtration, washed with water and air dried to afford the title compound (0.040 g, 0.12 mmol, 47 %).


$^1\text{H-NMR}$  (MeOD+Pyr)  $\delta$ : 7.44-7.34 (m, 2H); 6.75 (s, 1H), 6.67 (d, 1H,  $J = 7.9$  Hz); 6.38 (d, 1H,  $J = 8.1$  Hz); 3.43 (s, 3H); 2.48-2.31 (m, 3H), 1.99 (dd, 1H,  $J = 3.4$  and 3.5 Hz), 1.80-1.57 (m, 3H), 1.42 (t, 1H,  $J = 3.0$  Hz) ppm.

$^{13}\text{C-NMR}$  (MeOD+Pyr)  $\delta$ : 75.8, 175.5, 1721.8, 139.9, 139.6, 129.6, 125.6, 121.5, 118.1, 68.5, 42.6, 39.5, 38.6, 36.0, 35.3, 24.3, 21.6 ppm.

MS(ES) for  $\text{C}_{17}\text{H}_{20}\text{N}_2\text{O}_5$   $[\text{M}+\text{H}]^+$ : 333.0  $[\text{M}-\text{H}]^-$ : 331.10

### Example 81

Synthesis of (1S\*,2S\*,4R\*,5R\*,6S\*) 2-Amino-4-(m-trifluoromethyl)phenylcarbamoylmethyl bicyclo[3.1.0]hexane-2,6-dicarboxylic acid



A. 4(Z/E), (1S\*,2S\*,5R\*,6S\*) Diethyl 2-*tert*-butoxy-carbonylamino-4-(m-trifluoromethyl)phenylcarbamoylmethylene bicyclo[3.1.0]hexane-2,6-dicarboxylate.

-157-

A room temperature solution of the product of Example 74B (0.5 g, 1.25 mmol) in  $\text{CH}_2\text{Cl}_2$  was treated consecutively with oxalyl chloride (0.239 g, 1.89 mmol) and a catalytic amount of DMF. After the reaction mixture was stirred for one hour, the solvent was removed in vacuo. The crude product was reconstituted in  $\text{CH}_2\text{Cl}_2$  and then a solution of *m*-trifluoromethylaniline (0.402 g, 2.5 mmol) was added dropwise. The reaction mixture was allowed to stir at room temperature until the reaction was judged complete by TLC. The resulting mixture was concentrated to dryness and the crude product purified by filtration through a short pad of  $\text{SiO}_2$  to afford the title compounds (0.462 g, 0.85 mmol) 68 % as a mixture of diastereoisomers which were used in the next step without further separation.

B. (1S\*,2S\*,4R\*,5R\*,6S\*) Diethyl 2-*tert*-butyloxycarbonyl amino-4-(*m*-trifluoromethyl)phenylcarbamoyl methyl bicyclo[3.1.0]hexane-2,6-dicarboxylate.

A room temperature solution of the product of Example 81A (0.462 g, 0.85 mmol) and 10% Pd/C (0.023 g, 5 wt %) in 95% EtOH was vigorously stirred under a hydrogen atmosphere (1 atm) over night. The catalyst was removed via vacuum filtration through celite and the filtrate concentrated in vacuo to afford the title compound (0.347 g, 0.64 mmol, 70 %). MS(ES) for  $\text{C}_{26}\text{H}_{33}\text{F}_3\text{N}_2\text{O}_7$   $[\text{M}+\text{H}]^+$ : 543.10  $[\text{M}-\text{H}]^-$ : 541.23

C. (1S\*,2S\*,4R\*,5R\*,6S\*) Diethyl 2-Amino-4-(*m*-trifluoromethyl)phenylcarbamoyl methylbicyclo[3.1.0]hexane-2,6-dicarboxylate.

-158-


The title compound was prepared as in Example 74E to give the product in 56 % yield. MS(ES) for  $C_{21}H_{25}F_3N_2O_5$   $[M+H]^+$ : 443.2.

D. (1S\*,2S\*,4R\*,5R\*,6S\*) 2-Amino-4-(m-trifluoromethyl)phenylcarbamoylmethylbicyclo[3.1.0]hexane-2,6-dicarboxylic acid.

A solution of the product of Example 81C (0.140 g, 0.31 mmol) in THF was treated with 1N NaOH ( 0.8 mmol) and stirred at room temperature over night. The solvent was removed in vacuo and the crude product reconstituted in distilled water and acidified with 1N HCl. The product was isolated by isoelectric precipitation at pH = 3-5. The product was collected via vacuum filtration, washed with water and air dried to afford the title compound (0.110 g, 0.28mmol, 90 %). MS(ES) for  $C_{17}H_{17}F_3N_2O_5$   $[M+H]^+$ : 387.04.

### Example 82

Synthesis of (1S\*,2S\*,4R\*,5R\*,6S\*) 2-Amino-4-(p-methoxy)phenylcarbamoylmethyl bicyclo[3.1.0]hexane-2,6-dicarboxylic acid



A. 4(Z/E), (1S\*,2S\*,5R\*,6S\*) Diethyl 2-*tert*-butyloxy carbonylamino-4-(p-methoxy)phenylcarbamoylmethylene bicyclo[3.1.0]hexane-2,6-dicarboxylate.

A room temperature solution of the product of Example 74B (0.5 g, 1.25 mmol) in  $\text{CH}_2\text{Cl}_2$  was treated consecutively with oxalyl chloride (0.239 g, 1.89 mmol) and a catalytic amount of DMF. After the reaction mixture was stirred for one hour, the solvent was removed in vacuo. The crude product was reconstituted in  $\text{CH}_2\text{Cl}_2$  and then a solution of *p*-anisidine (0.307 g, 2.5 mmol) was added dropwise. The reaction mixture was allowed to stir at room temperature until the reaction was judged complete by TLC. The resulting mixture was concentrated to dryness and the crude product purified by filtration through a short pad of  $\text{SiO}_2$  to afford the title compounds (0.252 g, 0.5 mmol) 40 % as a mixture of diastereoisomers which were used in the next step without further separation.

B. (1S\*,2S\*,4R\*,5R\*,6S\*) Diethyl 2-tert-butylloxycarbonylamino-4-(*p*-methoxy)phenylcarbamoyl methylbicyclo[3.1.0]hexane-2,6-dicarboxylate.

A room temperature solution of the product of Example 82A (0.252 g, 0.5 mmol) and 10% Pd/C (0.013 g, 5 wt %) in 95% EtOH was vigorously stirred under a hydrogen atmosphere (1 atm) over night. The catalyst was removed via vacuum filtration through celite and the filtrate concentrated in vacuo to afford the title compound (0.164 g, 0.32 mmol, 65 %).

C. (1S\*,2S\*,4R\*,5R\*,6S\*) Diethyl 2-Amino-4-(*p*-methoxy)phenylcarbamoyl methylbicyclo[3.1.0]hexane-2,6-dicarboxylate.

The title compound was prepared as in Example 74E to give the product in 30 % yield.



-160-

$^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$ : 7.72 (s, 1H), 7.37 (d, 2H,  $J$  = 8.9 Hz), 6.81 (d, 2H,  $J$  = 8.9 Hz), 4.28-4.06 (dq, 4H), 3.77 (s, 3H), 3.68-2.72 (m, 1H), 2.49-2.44 (m, 2H), 2.34 (dd, 1H,  $J$  = 2.6 and 5.8 Hz), 2.15-1.96 (m, 4H), 1.68 (t, 1H,  $J$  = 3.0 Hz), 1.55 (dd, 1H,  $J$  = 8.1 and 14.0 Hz), 1.35-1.20 (dt, 6H) ppm

D. (1S\*,2S\*,4R\*,5R\*,6S\*) 2-Amino-4-(p-methoxy) phenylcarbamoylmethylbicyclo [3.1.0]hexane-2,6-dicarboxylic acid.


A solution of the product of Example 82C (0.025 g, 0.06 mmol) in THF was treated with 1N NaOH (0.12 mmol) and stirred at room temperature over night. The solvent was removed in vacuo and the crude product reconstituted in distilled water and acidified with 1N HCl. The product was isolated by isoelectric precipitation at pH = 3-5. The product was collected via vacuum filtration, washed with water and air dried to afford the title compound (0.008 g, 0.02mmol, 38 %)

$^1\text{H}$ - NMR ( $\text{D}_2\text{O}+\text{KOD}$ )  $\delta$ : 7.15 (d, 2H,  $J$  = 8.8 Hz); 6.82 (d, 2H,  $J$  = 8.9 Hz); 3.64 (s, 3H); 2.52-2.25 (m, 3H); 1.91-1.80 (m, 1H); 1.76 (d, 1H,  $J$  = 14.5 Hz), 1.38 (m, 1H), 1.30-1.24 (m, 2H) ppm.

$^{13}\text{C}$ -NMR ( $\text{D}_2\text{O}+\text{KOD}$ )  $\delta$ : 184.5, 182.4, 174.8, 130.7, 124.8, 114.9, 66.6, 56.1, 42.0, 41.4, 38.4, 35.9, 32.2, 25.5 ppm

Example 83

Synthesis of (1S\*,2S\*,4R\*,5R\*,6S\*) 2-Amino-4-(p-methyl)phenylcarbamoylmethyl bicyclo[3.1.0]hexane-2,6-dicarboxylic acid



A. 4(Z/E), (1S\*,2S\*,5R\*,6S\*) Diethyl 2-*tert*-butyloxy carbonylamino-4-(p-methyl)phenylcarbamoylmethylene bicyclo[3.1.0]hexane-2,6-dicarboxylate.

A room temperature solution of the product of Example 74B (0.5 g, 1.25 mmol) in CH<sub>2</sub>Cl<sub>2</sub> was treated consecutively with oxalyl chloride (0.239 g, 1.89 mmol) and a catalytic amount of DMF. After the reaction mixture was stirred for one hour, the solvent was removed in vacuo. The crude product was reconstituted in CH<sub>2</sub>Cl<sub>2</sub> and then a solution of *p*-toluidine (0.270 g, 2.5 mmol) was added dropwise. The reaction mixture was allowed to stir at room temperature until the reaction was judged complete by TLC. The resulting mixture was concentrated to dryness and the crude product purified by filtration through a short pad of SiO<sub>2</sub> to afford the title compounds (0.220 g, 0.45 mmol) 36 % as a mixture of diastereoisomers which were used in the next step without further separation.

-162-

B. (1S\*,2S\*,4R\*,5R\*,6S\*) Diethyl 2-tert-butyloxy carbonylamino-4-(p-methyl)phenylcarbamoylmethyl bicyclo[3.1.0]hexane-2,6-dicarboxylate.

A room temperature solution of the product of Example 83A (0.220 g, 0.45 mmol) and 10% Pd/C (0.011 g, 5 wt %) in 95% EtOH was vigorously stirred under an hydrogen atmosphere (1 atm) over night. The catalyst was removed via vacuum filtration through celite and the filtrate concentrated in vacuo to afford the title compound (0.137 g, 0.34 mmol, 65 %). MS(ES) for  $C_{26}H_{36}N_2O_7$   $[M+H]^+$ : 489.25  $[M-H]^-$ : 487.09.

C. (1S\*,2S\*,4R\*,5R\*,6S\*) Diethyl 2-Amino-4-(p-methyl) phenylcarbamoyl methylbicyclo [3.1.0]hexane-2,6-dicarboxylic acid.

The title compound was prepared as in Example 74E to give the product in 50 % yield.

$^1H$  NMR ( $CDCl_3$ ) $\delta$ : 8.04 (s, 1H), 7.41 (d, 1H, J= 8.4 Hz), 7.09 (d, 1H, J= 8.3 Hz), 4.26 (q, 2H, J= 7.1 Hz), 4.11 (q, 2H, J= 7.1 Hz), 2.86- 2.67 (m, 1H), 2.50-2.23 (m, 7H), 1.87-1.74 (m, 4H), 1.37-1.21 (dt, 6H) ppm.

D. (1S\*,2S\*,4R\*,5R\*,6S\*) 2-Amino-4-(p-methyl) phenylcarbamoylmethylbicyclo [3.1.0] hexane-2,6-dicarboxylic acid.

A solution of the product of Example 83C (0.055 g, 0.14 mmol) in THF was treated with 1N NaOH ( 0.28 mmol) and stirred at room temperature over night. The solvent was removed in vacuo and the crude product reconstituted in distilled water and acidified with 1N

-163-


HCl. The product was isolated by isoelectric precipitation at pH = 3-5. The product was collected via vacuum filtration, washed with water and air dried to afford the title compound (0.026 g, 0.07mmol, 55 %).

$^1\text{H-NMR}$  ( $\text{D}_2\text{O}+\text{KOD}$ )  $\delta$ : 7.26-7.14 (AA'BB', 4H), 2.58-2.30 (m, 3H), 2.24 (s, 3H); 2.02 (dd, 1H,  $J = 3.0$  and 6.1 Hz), 1.87 (d, 1H,  $J = 14.4$  Hz), 1.66 (dd, 1H,  $J = 2.9$  and 6.1 Hz), 1.46-1.19 (m, 2H) ppm.

$^{13}\text{C-NMR}$  ( $\text{D}_2\text{O}+\text{KOD}$ )  $\delta$ : 181.9, 174.3, 137.3, 135.7, 129.5, 122.3, 66.0, 41.7, 40.9, 37.8, 35.4, 31.8, 24.9, 19.9 ppm

#### Example 84

Synthesis of (1S\*,2S\*,4R\*,5R\*,6S\*) 2-Amino-4-(p-trifluoromethyl)phenylcarbamoylmethyl bicyclo[3.1.0]hexane-2,6-dicarboxylic acid



A. 4(Z/E), (1S\*,2S\*,5R\*,6S\*) Diethyl 2-*tert*-butyloxy carbonylamino-4-(p-trifluoromethyl)phenylcarbamoylmethylene bicyclo[3.1.0]hexane-2,6-dicarboxylate.

A room temperature solution of the product of Example 74B (0.5 g, 1.25 mmol) in  $\text{CH}_2\text{Cl}_2$  was treated consecutively with oxalyl chloride (0.239 g, 1.89 mmol) and a catalytic amount

-164-

of DMF. After the reaction mixture was stirred for one hour, the solvent was removed in vacuo. The crude product was reconstituted in  $\text{CH}_2\text{Cl}_2$  and then a solution of *p*-trifluoromethylaniline (0.290 g, 2.5 mmol) was added. The reaction mixture was allowed to stir at room temperature until the reaction was judged complete by TLC. The resulting mixture was concentrated to dryness and the crude product purified by filtration through a short pad of  $\text{SiO}_2$  to afford the title compound (0.340 g, 0.62 mmol) 50 % as a mixture of diastereoisomers which were used in the next step without further separation.

B. (1S\*,2S\*,4R\*,5R\*,6S\*) Diethyl 2-tert-butyloxy carbonylamino-4-(*p*-trifluoromethyl)phenylcarbamoylmethyl bicyclo[3.1.0]hexane-2,6-dicarboxylate.

A room temperature solution of the product of Example 84A (0.34 g, 0.62 mmol) and 10% Pd/C (0.017 g, 5 wt %) in 95% EtOH was vigorously stirred under an hydrogen atmosphere (1 atm) over night. The catalyst was removed via vacuum filtration through celite and the filtrate concentrated in vacuo to afford the title compound (0.320 g, 0.58 mmol, 93 %). MS(ES) for  $\text{C}_{26}\text{H}_{31}\text{F}_3\text{N}_2\text{O}_7$   $[\text{M}+\text{H}]^+$ : 545.24  $[\text{M}-\text{H}]^-$ : 543.27

C. (1S\*,2S\*,4R\*,5R\*,6S\*) Diethyl 2-Amino-4-(*p*-trifluoromethyl)phenylcarbamoyl methylbicyclo[3.1.0]hexane-2,6-dicarboxylate.

The title compound was prepared as in Example 74E to give the product in 54 % yield

$^1\text{H}$  NMR ( $\text{CDCl}_3$ ) $\delta$ : 8.50 (s, 1H), 7.797.51(m, 4H), 4.34-4.01 (m, 4H), 2.97- 2.17 (m, 6H), 1.98-1.65 (m, 4H), 1.38-1.18

-165-

(m, 6H) ppm. MS(ES) for  $C_{21}H_{25}F_3N_2O_5$   $[M+H]^+$ : 443.01;  $[M-H]^-$ : 441.15

D. (1S\*,2S\*,4R\*,5R\*,6S\*) 2-Amino-4-(p-trifluoromethyl)phenylcarbamoylmethylbicyclo [3.1.0]hexane-2,6-dicarboxylic acid.


A solution of the product of Example 84C (0.067 g, 0.15 mmol) in THF was treated with 1N NaOH (0.30 mmol) and stirred at room temperature over night. The solvent was removed in vacuo and the crude product reconstituted in distilled water and acidified with 1N HCl. The product was isolated by isoelectric precipitation at pH = 3-5. The product was collected via vacuum filtration, washed with water and air dried to afford the title compound (0.033 g, 0.08mmol, 57 %).

$^1H$ -NMR ( $D_2O$ +KOD)  $\delta$ : 7.65 (d, 2H, J = 8.6 Hz); 7.54 (d, 2H, J = 8.8 Hz); 2.65-2.40 (m, 3H); 2.04 (d, 1H, J = 3.1 and 6.2 Hz); ); 1.89 (d, 1H, J = 14.4 Hz), 1.69 (d, 1H, J = 2.9 and 5.9 Hz); 1.48-1.35 (m, 2H) ppm. MS(ES) for  $C_{17}H_{17}F_3N_2O_5$   $[M+H]^+$ : 387.04  $[M-H]^-$ : 385.07

#### Example 85

Synthesis of (1S\*,2S\*,4R\*,5R\*,6S\*) 2-Amino-4-(m-chloro)phenylcarbamoylmethyl bicyclo[3.1.0]hexane-2,6-dicarboxylic acid

-166-



A. 4(Z/E), (1S\*,2S\*,5R\*,6S\*) Diethyl 2-*tert*-butoxy carbonylamino-4-(*m*-chloro)phenylcarbamoylmethylene bicyclo[3.1.0]hexane-2,6-dicarboxylate.

A room temperature solution of the product of Example 74B (0.5 g, 1.25 mmol) in CH<sub>2</sub>Cl<sub>2</sub> was treated consecutively with oxalyl chloride (0.239 g, 1.89 mmol) and a catalytic amount of DMF. After the reaction mixture was stirred for one hour, the solvent was removed in vacuo. The crude product was reconstituted in CH<sub>2</sub>Cl<sub>2</sub> and then a solution of *m*-chloroaniline (0.290 g, 2.5 mmol) was added dropwise. The reaction mixture was allowed to stir at room temperature until the reaction was judged complete by TLC. The resulting mixture was concentrated to dryness and the crude product purified by filtration through a short pad of SiO<sub>2</sub> to afford the title compound (0.340 g, 0.62 mmol) 50 % as a mixture of diastereoisomers which were used in the next step without further separation.

B. (1S\*,2S\*,4R\*,5R\*,6S\*) Diethyl 2-*tert*-butoxy carbonylamino-4-(*m*-chloro)phenylcarbamoylmethyl bicyclo[3.1.0]hexane-2,6-dicarboxylate.

A room temperature solution of the product of Example 85A (0.34 g, 0.62 mmol) and 10% Pd/C (0.017 g, 5 wt %) in 95% EtOH was vigorously stirred under a hydrogen atmosphere (1 atm) over night. The catalyst was removed via vacuum

-167-

filtration through celite and the filtrate concentrated in vacuo to afford the title compound.

C. (1S\*,2S\*,4R\*,5R\*,6S\*) Diethyl 2-Amino-4-(m-chloro)phenylcarbamoyl methylbicyclo[3.1.0]hexane-2,6-dicarboxylic acid.

The title compound was prepared as in Example 74E to give the product.

$^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$ : 8.27 (bs, 1H), 7.52 (m, 2H), 7.27 (m, 2H), 7.04 (m, 1H), 4.22 (q, 2H,  $J = 7.1$  Hz), 4.08 (q, 2H), 2.89-2.72 (m, 1H), 2.51-2.28 (m, 5H), 1.85-1.73 (m, 4H), 1.25 (dt, 6H) ppm.

MS(ES) for  $\text{C}_{20}\text{H}_{25}\text{ClN}_2\text{O}_5$   $[\text{M}+\text{H}]^+/-\text{Cl}$ : 375.03;  $[\text{M}-\text{H}]^-/-\text{Cl}$ : 373.15

D. (1S\*,2S\*,4R\*,5R\*,6S\*) 2-Amino-4-(m-chloro)phenyl carbamoylmethylbicyclo [3.1.0]hexane-2,6-dicarboxylic acid.

A solution of the product of Example 85C (0.067 g, 0.15 mmol) in THF was treated with 1N NaOH (0.30 mmol) and stirred at room temperature over night. The solvent was removed in vacuo and the crude product reconstituted in distilled water and acidified with 1N HCl. The product was isolated by isoelectric precipitation at pH = 3-5. The product was collected via vacuum filtration, washed with water and air dried to afford the title compound


MS(ES) for  $\text{C}_{16}\text{H}_{17}\text{ClN}_2\text{O}_5$   $[\text{M}+\text{H}]^+/-\text{Cl}$ : 319.18  $[\text{M}-\text{H}]^-/-\text{Cl}$ : 317.23

$^{13}\text{C}$ -NMR ( $\text{D}_2\text{O}+\text{Pyr}$ )  $\delta$ : 178.5, 175.2, 171.8, 136.7, 136.2, 128.0, 123.7, 123.1, 119.7, 66.8, 59.33, 40.6, 37.4, 37.2, 32.6, 32.2, 25.1 ppm.



Example 86

Synthesis of (1S\*,2S\*,4R\*,5R\*,6S\*) 2-Amino-4-carboxymethyl bicyclo[3.1.0]hexane-2,6-dicarboxylic acid



A. 4(Z/E), (1S\*,2S\*,5R\*,6S\*) Diethyl 2-tert-butyloxy carbonylamino-4-methoxycarbonylmethylene bicyclo[3.1.0]hexane-2,6-dicarboxylate.

A room temperature solution of the product of Example 74B (0.5 g, 1.25 mmol) in CH<sub>2</sub>Cl<sub>2</sub> was treated consecutively with oxalyl chloride (0.239 g, 1.89 mmol) and a catalytic amount of DMF. After the reaction mixture was stirred for one hour, the solvent was removed in vacuo. The crude product was reconstituted in CH<sub>2</sub>Cl<sub>2</sub> and then a solution of trifluoromethylaniline was added. The reaction mixture was allowed to stir at room temperature until the reaction was judged complete by TLC. The resulting mixture was concentrated to dryness and the crude product purified by filtration through a short pad of SiO<sub>2</sub> to afford the title

compound as a mixture of diastereoisomers which were used in the next step without further separation.

B. (1S\*,2S\*,4R\*,5R\*,6S\*) Diethyl 2-tert-butyloxy carbonylamino-4-methoxycarbonylmethyl bicyclo[3.1.0]hexane-2,6-dicarboxylate.

A room temperature solution of the product described above (2 g, 5.0 mmol) and 10% Pd/C (0.250 g, 5 wt %) in 95% EtOH was vigorous stirred under an hydrogen atmosphere (1 atm) over night. The catalyst was removed via vacuum filtration through celite and the filtrate concentrated in vacuo to afford the title compound (1.7 g, 4.2 mmol, 84 %). MS(ES) for  $C_{10}H_{13}NO_6$   $[M+H]^+$ : 244.16  $[M-H]^-$ : 242.19.

C. (1S\*,2S\*,4R\*,5R\*,6S\*) Diethyl 2-Amino-4-methoxycarbonyl methylbicyclo [3.1.0] hexane-2,6-dicarboxylic acid.

The title compound was prepared as example 74E to give the product in 70 %  $^1H$  NMR ( $CDCl_3$ ) $\delta$ : 4.23-4.00 (dq, 4H), 3.60 (s, 3H), 2.70-2.60 (m, 1H), 2.51-2.33 (m, 2H), 2.26 (dd, 1H,  $J = 3.0$  and  $6.2$  Hz), 2.07 (d, 1H,  $J = 14.8$  Hz), 1.83 (dd, 1H,  $J = 3.0$  and  $6.1$  Hz), 1.74 (brs, 2H), 1.67 (t, 1H,  $J = 3.0$  Hz), 1.43 (dd, 1H,  $J = 8.3$  and  $14.8$  Hz), 1.30 1.15 (m, 6H). ppm

D. (1S\*,2S\*,4R\*,5R\*,6S\*) 2-Amino-4-carboxymethyl bicyclo[3.1.0]hexane-2,6-dicarboxylic acid.

A solution of the product of Example 86C (0.180 g, 0.6 mmol) in THF was treated with 1N NaOH ( 1.2 mmol) and stirred at room temperature over night. The solvent was removed in vacuo and the crude product reconstituted in distilled water and acidified with 1N HCl. The product was isolated by ion


-170-

exchange chromatography, washed with water and air dried to afford the title compound (0.052 g, 0.21mmol, 40 %).

$^1\text{H}$ - NMR ( $\text{D}_2\text{O}$  + Pyr) $\delta$ : 2.89-2.37 (m, 3H), 2.08 (dd, 1H,  $J$  = 3.1 and 6.1 Hz), 1.93-1.62 (m, 3H), 1.50 (t, 1H,  $J$  = 3.1 Hz) ppm.  $^{13}\text{C}$ - NMR ( $\text{D}_2\text{O}$  + Pyr) $\delta$ : 183.9, 182.6, 179.0, 70.4, 45.1, 41.5, 41.2, 36.4, 36.0, 28.7 ppm. MS(ES) for  $\text{C}_{10}\text{H}_{13}\text{NO}_6$   $[\text{M}+\text{H}]^+$ : 244.16  $[\text{M}-\text{H}]^-$ : 242.19.

### Example 87

Synthesis of (1S\*,2S\*,4R\*,5R\*,6S\*) 2-Amino-4-benzoxazol-2-yl-methylbicyclo[3.1.0]hexane-2,6-dicarboxylic acid



A. 4(Z/E),1S\*,2S\*,5R\*,6S\* Diethyl 2-tert-butyloxy-carbonylamino-4-(benzoxazol-2-yl)methylene bicyclo[3.1.0]-hexane-2,6-dicarboxylate.

To a 0 °C solution of diethyl benzoxazol-2-ylmethylphosphonate (previously prepared) 0.454 1.68 mmol in anhydrous THF, was added 1M NaHMDS (1.68 mmol) by dropwise addition. The resulting reaction mixture was stirred at 0 °C for 1 h, after which time a solution of (1S\*,2S\*,5R\*,6R\*) diethyl2-N-tert-butyloxycarbonylamino-4-oxobicyclo[3.1.0]hexane-2,6-dicarboxylate (previously disclosed US 5,958,960 example 5a) in THF was added dropwise. The reaction mixture

-171-

was allowed to room temperature as it stirred 4 h. The reaction was quenched with saturated aqueous  $\text{NH}_4\text{Cl}$  solution and the product extracted with ethyl acetate. The combined organic extracts were dried over sodium sulfate, concentrated to dryness in vacuo, and purified by  $\text{SiO}_2$  chromatography. ( EtOAc/ hexanes. 3:7) to afford 0.380 g (1.12 mmol, 72 %) of the title compound as a (1:1.5) mixture of diastereoisomers 1:1.5 which were used in the next step without further separation. MS(ES) for  $\text{C}_{25}\text{H}_{30}\text{N}_2\text{O}_7$   $[\text{M}+\text{H}]^+$ : 471.3.

B. (1S\*,2S\*,4R\*,5R\*,6S\*) Diethyl 2-tert-butyloxycarbonyl-amino-4-(benzoxazol-2-yl)methylbicyclo[3.1.0]hexane-2,6-dicarboxylate.

A room temperature solution of the product of 87A (0.380 g, 0.8 mmol) and 10% Pd/C (0.020 g, 5 wt %) in 95% EtOH was vigorously stirred under an hydrogen atmosphere (1 atm) over night. The catalyst was removed via vacuum filtration through celite and the filtrate concentrated in vacuo to afford the title compound (0.275 g, 0.58 mmol, 72 %). MS(ES) for  $\text{C}_{25}\text{H}_{32}\text{N}_2\text{O}_7$   $[\text{M}+\text{H}]^+$ : 473.07.

C. (1S\*,2S\*,4R\*,5R\*,6S\*) Diethyl 2-Amino-4-(benzoxazol-2-yl)methylbicyclo [3.1.0]hexane-2,6-dicarboxylic acid.

The title compound was prepared as example 74E to give the product in 53 %.

$^1\text{H}$  NMR ( $\text{CDCl}_3$ ) $\delta$ : 7.66-7.62 (m, 1H), 7.47-7.43 (m, 1H), 7.31-7.24 (m, 2H), 4.26 (q, 2H,  $J$  = 7.1 Hz), 4.08 (q, 2H,  $J$  = 7.1 Hz), 3.09-2.90 (m, 3H), 2.37 (dd, 1H,  $J$  = 2.9 and 6.0 Hz), 2.25 (d, 1H,  $J$  = 14.8 Hz), 2.01 (d, 1H,  $J$  = 3.1 and 5.9 Hz),

-172-

1.79-1.74 (m, 3H), 1.53 (dd, 1H,  $J = 7.9$  and  $14.8$  Hz), 1.33 (t, 3H,  $J = 7.1$  Hz), 1.22 (t, 3H,  $J = 7.1$  Hz) ppm. MS(ES) for  $C_{20}H_{24}N_2O_7$   $[M+H]^+$ : 373.18.


D. (1S\*,2S\*,4R\*,5R\*,6S\*) 2-Amino-4-benzoxazol-2-yl-methylbicyclo[3.1.0]hexane-2,6-dicarboxylic acid

A solution of the product of Example 87C (0.080 g, 0.21 mmol) in THF was treated with 1N NaOH (0.43 mmol) and stirred at room temperature over night. The solvent was removed in vacuo and the crude product reconstituted in distilled water and acidified with 1N HCl. The product was isolated by isoelectric precipitation at pH = 3-5. The product was collected via vacuum filtration, washed with water and air dried to afford the title compound (0.040 g, 0.12mmol, 59 %).

$^1H$ - NMR ( $D_2O + KOD$ ) $\delta$ : 7.60-7.52 (m, 2H), 7.33-7.29 (m, 2H), 3.04-2.98 (m, 2H), 2.68-2.56 (m, 1H), 2.04 (dd, 1H,  $J = 3.2$  and  $6.4$  Hz), 1.92 (d, 1H,  $J = 14.6$  Hz), 1.74-1.71 (m, 1H), 1.48-1.45 (m, 2H) ppm.  $^1H$ - NMR ( $D_2O + KOD$ ) $\delta$ : 183.3, 181.2, 167.6, 125.0, 124.5, 118.5, 110.8, 66.1, 41.1, 38.6, 35.7, 33.2, 32.0, 25.0 ppm. MS(ES) for  $C_{16}H_{16}N_2O_5$   $[M+H]^+$ : 317.02.

#### Example 88

Synthesis of (1S\*,2S\*,4R\*,5R\*,6S\*) 2-Amino-4-benzotiazol-2-yl-methylbicyclo[3.1.0]hexane-2,6-dicarboxylic acid



-173-

A. 4(Z/E), (1*S*\*, 2*S*\*, 5*R*\*, 6*S*\*) Diethyl 2-*tert*-butyloxycarbonylamino-4-(benzotiazol-2-yl)methylene bicyclo[3.1.0]hexane-2,6-dicarboxylate.

To a 0 °C solution of diethyl benzotiazol-2-yl-methylphosphonate (previously prepared) 0.360g, 1.26 mmol in anhydrous THF, was added 1M NaHMDS (1.26 mmol) by dropwise addition. The resulting reaction mixture was stirred at 0 °C for 1 h, after which time a solution of (1*S*\*, 2*S*\*, 5*R*\*, 6*R*\*) diethyl-2-*N*-*tert*-butyloxycarbonylamino-4-oxobicyclo[3.1.0]hexane-2,6-dicarboxylate (previously disclosed US 5,958,960 example 5a) 0.3g, 0.84 mmol in THF was added dropwise. The reaction mixture was allowed to room temperature as it stirred 4 h. The reaction was quenched with saturated aqueous NH<sub>4</sub>Cl solution and the product extracted with ethyl acetate. The combined organic extracts were dried over sodium sulfate, concentrated to dryness in vacuo, and purified by SiO<sub>2</sub> chromatography. (EtOAc/ hexanes. 3:7) to afford 0.30 g (0.61 mmol, 72 %) of the title compound as a (1:1.5) mixture of diastereoisomers which were used in the next step without further separation. MS(ES) for C<sub>25</sub>H<sub>30</sub>N<sub>2</sub>O<sub>6</sub>S [M+H]<sup>+</sup>: 486.98 [M-H]: 485.08.

B. (1*S*\*, 2*S*\*, 4*R*\*, 5*R*\*, 6*S*\*) Diethyl 2-*tert*-butyloxycarbonylamino-4-(benzotiazol-2-yl)methyl bicyclo[3.1.0]hexane-2,6-dicarboxylate.

A room temperature solution of the product of 88A (0.30 g, 0.61 mmol) and 10% Pd/C (0.016 g, 5 wt %) in 95% EtOH was vigorously stirred under an hydrogen atmosphere (1 atm) over night. The catalyst was removed via vacuum filtration

-174-

through celite and the filtrate concentrated in vacuo to afford the title compound (0.250 g, 0.51 mmol, 61 %). MS(ES) for  $C_{25}H_{32}N_2O_6S$   $[M+H]^+$ : 489.05.

C. (1*S*\*,2*S*\*,4*R*\*,5*R*\*,6*S*\*) Diethyl 2-Amino-4-(benzotiazol-2-yl)methylbicyclo [3.1.0] hexane-2,6-dicarboxylic acid.

The title compound was prepared as example 74E to give the product in 90 %.

$^1H$  NMR ( $CDCl_3$ ) $\delta$ : 7.95 (ddd, 1H,  $J$  = 7.8, 1.2 and 0.5 Hz), 7.80 (ddd, 1H), 7.47-7.25 (m, 2H), 4.28 (q, 2H,  $J$  = 7.1 Hz), 4.08 (q, 2H,  $J$  = 7.1 Hz), 3.31-3.09 (m, 2H), 2.9 (m, 1H), 2.39 (dd, 1H,  $J$  = 2.9 and 6.2 Hz), 2.33 (d, 1H,  $J$  = 14.7 Hz), 2.01 (d, 1H,  $J$  = 3.0 and 5.9 Hz), 1.76-1.73 (m, 3H), 1.49 (dd, 1H,  $J$  = 8.3 and 14.7 Hz), 1.34 (t, 3H,  $J$  = 7.1 Hz), 1.22 (t, 3H,  $J$  = 7.1 Hz) ppm.

D. 1*S*\*,2*S*\*,4*R*\*,5*R*\*,6*S*\*) 2-Amino-4-benzoxazol-2-yl-methylbicyclo[3.1.0]hexane-2,6-dicarboxylic acid

A solution of the product of Example 88C (0.080 g, 0.20 mmol) in THF was treated with 1N NaOH (0.40 mmol) and stirred at room temperature over night. The solvent was removed in vacuo and the crude product reconstituted in distilled water and acidified with 1N HCl. The product was isolated by isoelectric precipitation at pH = 3-5. The product was collected via vacuum filtration, washed with water and air dried to afford the title compound (0.048 g, 0.14mmol, 70 %).


$^1H$ - NMR ( $D_2O$  + KOD) $\delta$ : 7.87-7.77 (m, 2H), 7.46-7.29 (m, 2H), 3.21-2.99 (m, 2H), 2.67-2.56 (m, 1H), 2.03 (dd, 1H,  $J$  = 3.2

-175-

and 6.4 Hz), 1.90 (d, 1H,  $J = 14.6$  Hz), 1.72-1.69 (m, 1H), 1.43-1.30 (m, 2H) ppm.  $^{13}\text{C}$ - NMR ( $\text{D}_2\text{O} + \text{KOD}$ )  $\delta$ : 181.8, 180.4, 172.2, 150.3, 132.9, 124.9, 123.7, 120.6, 119.8, 64.6, 39.8, 39.4, 36.8, 34.2, 30.5, 23.6 ppm. MS(ES) for  $\text{C}_{16}\text{H}_{16}\text{N}_2\text{O}_4\text{S}$   $[\text{M}+\text{H}]^+$ : 332.94  $[\text{M}-\text{H}]^-$ : 331.03.

### Example 89

Synthesis of (1S\*,2S\*,4R\*,5R\*,6S\*) 2-Amino-4-benzylcarbamoyl methylbicyclo[3.1.0]hexane-2,6-dicarboxylic acid



A. 4(Z/E), (1S\*,2S\*,5R\*,6S\*) Diethyl 2-tert-butyloxy-carbonylamino-4-benzylcarbamoylmethylenebicyclo[3.1.0]-hexane-2,6-dicarboxylate.

A room temperature solution of the product of Example 74B (0.5 g, 1.25 mmol) in  $\text{CH}_2\text{Cl}_2$  was treated consecutively with oxalyl chloride (0.239 g, 1.89 mmol) and a catalytic amount of DMF. After the reaction mixture was stirred for one hour, the solvent was removed in vacuo. The crude product was reconstituted in  $\text{CH}_2\text{Cl}_2$  and then a solution of benzylamine (0.267 g, 2.5 mmol) was added dropwise. The reaction mixture was allowed to stir at room temperature until the reaction was judged complete by TLC. The resulting mixture was concentrated to dryness and the crude product purified by filtration through a short pad of  $\text{SiO}_2$  to afford the title compound (0.350 g, 0.78 mmol) 62 % as a mixture of



-176-

diastereoisomers which were used in the next step without further separation. MS(ES) for  $C_{26}H_{34}N_2O_7$   $[M+H]^+$ : 487.24  $[M-H]^-$ : 485.29.

B. (1S\*,2S\*,4R\*,5R\*,6S\*) Diethyl 2-tert-butyloxy-carbonylamino-4-benzylcarbamoyl methylbicyclo[3.1.0]hexane-2,6-dicarboxylate.

A room temperature solution of the product of 89A (0.20 g, 0.41 mmol) and 10% Pd/C (0.015 g, 5 wt %) in 95% EtOH was vigorous stirred under an hydrogen atmosphere (1 atm) over night. The catalyst was removed via vacuum filtration through celite and the filtrate concentrated in vacuo to afford the title compound (0.150 g, 0.30 mmol, 75 %). MS(ES) for  $C_{26}H_{36}N_2O_7$   $[M+H]^+$ :

C. (1S\*,2S\*,4R\*,5R\*,6S\*) Diethyl 2-Amino-4-benzylcarbamoyl methylbicyclo [3.1.0]hexane-2,6-dicarboxylic acid  
The title compound was prepared as example 74E to give the product.

D. (1S\*,2S\*,4R\*,5R\*,6S\*) 2-Amino-4-benzylcarbamoylmethyl-bicyclo [3.1.0]hexane-2,6-dicarboxylic acid.

A solution of the product of Example 89C ( 0.150 g, 0.3 mmol) in THF was treated with 1N NaOH ( 0.6 mmol) and stirred at room temperature over night. The solvent was removed in vacuo and the crude product reconstituted in distilled water and acidified with 1N HCl. The product was isolated by isoelectric precipitation at pH = 3-5. The product was collected via vacuum filtration, washed with water and air dried to afford the title compound ( 0.055g, 0.16 mmol, 54 %).


-177-

$^1\text{H}$ - NMR ( $\text{D}_2\text{O}$  + KOD)  $\delta$ : 7.33-7.24 (m, 5H), 4.69 (s, 2H), 2.55-2.21 (m, 3H, 2.99 (dd, 1H,  $J$  = 3.2 and 5.9 Hz), 1.80 (d, 1H,  $J$  = 14.4 Hz), 1.60 (dd, 1H,  $J$  = 2.7 and 6.5 Hz), 1.43-1.32 (m, 2H) ppm.  $^{13}\text{C}$ - NMR ( $\text{D}_2\text{O}$  + KOD)  $\delta$ : 185.2, 183.6, 177.2, 139.9, 130.6, 129.2, 129.0, 44.7, 42.9, 42.9, 39.7, 37.3, 33.6 ppm

-178-

We claim:

1. A compound of the formula




wherein:

X is CH<sub>2</sub>, O, or NH;

Y is O, S, N or H;

A is a bond, O, N, (1-10C) alkyl, (2-10C) alkenyl or (2-10C) alkynyl;

R is hydrogen, (1-10C) alkyl, (2-10C) alkenyl, (3-6C) alkynyl, aryl, heterocyclyl or substituted aryl;  
or the group XC(Y)AR is



where Q is O, S or NH;

or a pharmaceutically acceptable metabolically labile ester or amide thereof;

or a pharmaceutically acceptable salt thereof.

2. The compound (or salt thereof) of Claim 1 wherein (1-10C) alkyl is methyl, ethyl, n-propyl, isopropyl, n-butyl, t-butyl, n-pentyl, isopentyl, n-hexyl, heptyl, n-octyl, nonyl or decyl; (2-10C) alkenyl is allyl, allenyl, 1-butenyl, 1-pentenyl, 3-nonenyl or 5-decenyl; (2-6C) alkynyl is ethynyl, propynyl, butynyl or pentynyl; aryl is phenyl, substituted phenyl or naphthyl; and arylalkyl is benzyl, 2-nitro benzyl, or 1-phenylethyl.

3. The compound (or salt thereof) of any one of Claims 1 or 2 wherein:

R is 2-methoxyphenyl, 3-methoxyphenyl, 4-methoxyphenyl, 3-methylphenyl, 4-methylphenyl, 2-chlorophenyl, 3-chlorophenyl, 4-chlorophenyl, 3-fluorophenyl, 2-fluorophenyl, 4-fluorophenyl, 2,6-dichlorophenyl, 2,4-dichlorophenyl, 3,5-dichlorophenyl, 3,4-dichlorophenyl, methyl, 2-naphthyl, 1-naphthyl, 3-methylphenyl, phenyl, thiophenyl, 3-trifluoromethylphenyl, 2,3-dichlorophenyl, 1H-indol-3-yl cyclopropanyl, 1H-indol-2-yl, 4-hydroxyphenyl, 3-hydroxyphenyl, 2-hydroxyphenyl, 4-biphenyl, 1-isoquinolyl, 3-pyridinyl, 2-pyridinyl, 3,5-difluorophenyl, 4-pyridinyl, 2-methylphenyl, 2-quinoxaliny, hydrogen, 3-carboxyphenyl, 1H-4-imidazolyl, 2-carboxyphenyl, 4-carboxyphenyl, 2-trifluoromethylphenyl, benzyl, 4-trifluoromethylphenyl;

A is a bond, methyl, O, NH, ethenyl or ethynyl;

Y is O, S or H,H; and

X is CH<sub>2</sub>, O or NH.

4. The compound (or salt thereof) of any one of claims 1-3 wherein R is heterocyclyl or substituted aryl.

5. The compound (or salt thereof) of any one of claims 1-4 wherein R is 1H-indolyl, 2-naphthyl, 3-chlorophenyl or 2-methoxyphenyl.

6. The compound (or salt thereof) of any one of claims 1-5 wherein Y is O or S.

7. The compound (or salt thereof) of any one of claims 1-6 wherein A is CH<sub>2</sub> or a bond.

8. The compound of formula I which is (1S\*, 2S\*, 4S\*, 5R\*, 6S\*) 2-amino-4-[3-

3-chlorophenyl)ureido]bicyclo[3.1.0]hexane-2,6-dicarboxylic acid.

9. A method for affecting the cAMP-linked metabotropic glutamate receptors in a patient, which comprises administering to a patient requiring modulated excitatory amino acid neurotransmission a pharmaceutically-effective amount of a compound of Claim 1.

10. A method for affecting the cAMP-linked metabotropic glutamate receptors in a patient, which comprises administering to a patient requiring modulated excitatory amino acid neurotransmission a pharmaceutically-effective amount of the compound of Claim 8.

11. A method for treating a neurological disorder in a patient which comprises administering to the patient in need of treatment thereof a pharmaceutically-effective amount of a compound of Claim 1.

12. The method of Claim 11 wherein said neurological disorder is cerebral deficits subsequent to cardiac bypass and grafting; cerebral ischemia; spinal cord trauma; head trauma; Alzheimer's Disease; Huntington's Chorea; amyotrophic lateral sclerosis; AIDS-induced dementia; perinatal hypoxia; hypoglycemic neuronal damage; ocular damage and retinopathy; cognitive disorders; idiopathic and drug-induced Parkinson's Disease; muscular spasms; migraine headaches; urinary incontinence; drug tolerance, withdrawal, and cessation; smoking cessation; emesis; brain edema; chronic pain; sleep disorders; convulsions; Tourette's syndrome; attention deficit disorder; and tardive dyskinesia.

-181-

13. The method of Claim 12 wherein said neurological disorder is drug tolerance, withdrawal, and cessation; or smoking cessation.

14. A method for treating a neurological disorder in a patient which comprises administering to the patient in need of treatment thereof a pharmaceutically-effective amount of a compound of Claim 8.

15. The method of Claim 14 wherein said neurological disorder is cerebral deficits subsequent to cardiac bypass and grafting; cerebral ischemia; spinal cord trauma; head trauma; Alzheimer's Disease; Huntington's Chorea; amyotrophic lateral sclerosis; AIDS-induced dementia; perinatal hypoxia; hypoglycemic neuronal damage; ocular damage and retinopathy; cognitive disorders; idiopathic and drug-induced Parkinson's Disease; muscular spasms; migraine headaches; urinary incontinence; drug tolerance, withdrawal, and cessation; smoking cessation; emesis; brain edema; chronic pain; sleep disorders; convulsions; Tourette's syndrome; attention deficit disorder; and tardive dyskinesia.

16. The method of Claim 15 wherein said neurological disorder is drug tolerance, withdrawal, and cessation; or smoking cessation.

17. A method for treating a psychiatric disorder in a patient which comprises administering to the patient in need of treatment thereof a pharmaceutically-effective amount of a compound of Claim 1.

18. The method of Claim 17 wherein said psychiatric disorder is schizophrenia, anxiety and related disorders, depression, bipolar disorders, psychosis, and obsessive compulsive disorders.

19. The method of Claim 18 wherein said psychiatric disorder is anxiety and related disorders.

20. A method for treating a psychiatric disorder in a patient which comprises administering to the patient in need of treatment thereof a pharmaceutically-effective amount of a compound of Claim 8.

21. The method of Claim 20 wherein said psychiatric disorder is schizophrenia, anxiety and related disorders, depression, bipolar disorders, psychosis, and obsessive compulsive disorders.

22. The method of Claim 21 wherein said psychiatric disorder is anxiety and related disorders.

23. A pharmaceutical formulation comprising a compound of Claim 1 in combination with one or more pharmaceutically-acceptable carriers, diluents, or excipients.

24. A pharmaceutical formulation comprising the compound of Claim 8 in combination with one or more pharmaceutically-acceptable carriers, diluents, or excipients.

25. A novel compound of formula I substantially as hereinbefore described with reference to any of the Examples.

26. A method for affecting the CAMP-linked metabotropic glutamate receptors in a patient, which comprises administering to a patient requiring modulated excitatory amino acid neurotransmission a pharmaceutically effective amount of a compound of formula I substantially as

-183-

hereinbefore described with reference to any of the Examples.

27. A process for preparing a novel compound of formula I substantially as hereinbefore described with reference to any of the Examples.



# INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 02/01247

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7	C07C275/18	C07C275/34	C07C335/16	C07D333/38	C07C233/52
	C07C233/81	C07C237/38	C07C235/54	C07C233/63	C07C235/62
	C07D209/42	C07D217/26	C07D213/81	C07D213/82	C07D241/44

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07C C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, CHEM ABS Data

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 98 51655 A (LILLY CO ELI ;MASSEY STEVEN MARC (US); MONN JAMES ALLEN (US); VALL) 19 November 1998 (1998-11-19) cited in the application page 29 -page 68 -----	1-27

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

\* Special categories of cited documents :

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \* & \* document member of the same patent family

Date of the actual completion of the international search

23 May 2002

Date of mailing of the international search report

03/06/2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Bader, K

# INTERNATIONAL SEARCH REPORT

International Application No.

PCT/US 02/01247

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07C235/84 C07D233/90 C07C229/48 C07C271/38 C07C237/14  
 C07D263/56 C07D277/64 A61K31/00 A61P25/18 A61P25/30  
 A61P13/04 A61P25/00

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

° Special categories of cited documents :

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \*G\* document member of the same patent family

Date of the actual completion of the international search

23 May 2002

Date of mailing of the international search report

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
 NL - 2280 HV Rijswijk  
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
 Fax: (+31-70) 340-3016

Authorized officer

Bader, K

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 02/01247

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9851655	A	19-11-1998	AT 198190 T 15-01-2001
			AU 729632 B2 08-02-2001
			AU 7386998 A 08-12-1998
			BR 9808787 A 11-07-2000
			CN 1255117 T 31-05-2000
			DE 69800430 D1 25-01-2001
			DE 69800430 T2 13-06-2001
			DK 878463 T3 19-02-2001
			EP 0878463 A1 18-11-1998
			ES 2152738 T3 01-02-2001
			HU 0001404 A2 28-08-2000
			JP 2001525825 T 11-12-2001
			NO 995480 A 09-11-1999
			NZ 500954 A 29-06-2001
			PL 336793 A1 17-07-2000
			PT 878463 T 30-04-2001
			SI 878463 T1 30-06-2001
			TR 9902777 T2 21-07-2000
			US 5958960 A 28-09-1999
			WO 9851655 A1 19-11-1998
			US 6160009 A 12-12-2000
			US 6268507 B1 31-07-2001
			ZA 9803930 A 08-11-1999